

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


As rescanning documents *will not* correct images,
Please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification n^o 7 : C07D 471/06, 487/06, A61K 31/4375, 31/55, 31/551, A61P 35/00, C07D 519/00 // (C07D 471/06, 221:00, 209:00) (C07D 487/06, 223:00, 209:00) (C07D 487/06, 243:00, 209:00) (C07D 519/00, 487:00, 487:00)</p>	A1	<p>(11) International Publication Number: WO 00/42040</p> <p>(43) International Publication Date : 20 July 2000 (20.07.00)</p>
<p>(21) International Application Number: PCT/US00/00411</p> <p>(22) International Filing Date: 10 January 2000 (10.01.00)</p> <p>(30) Priority Data: 60/115,431 11 January 1999 (11.01.99) US</p> <p>(71) Applicants (for all designated States except US): AGOURON PHARMACEUTICALS, INC. [US/US]; 10350 North Torrey Pines Road, La Jolla, CA 92037-1020 (US). CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): WEBBER, Stephen, Evan [US/US]; 3531 Millikin Avenue, San Diego, CA 92122 (US). CANAN-KOCH, Stacie, S. [US/US]; 5940 La Jolla Mesa Drive, La Jolla, CA 92037 (US). TIKHE, Jayashree [IN/US]; 9665 Genesee Avenue #D-2, San Diego, CA 92121 (US). THORESEN, Lars, Henrik [US/US]; Texas A</p>	<p>& M University, Department of Chemistry, P.O. Box 30012, College Station, TX 77482-3012 (US).</p> <p>(74) Agents: HERBERT, Toni-Junell; Shanks & Herbert, TransPotomac Plaza, Suite 306, 1033 N. Fairfax Street, Alexandria, VA 22314 (US) et al.</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	

(54) Title: TRICYCLIC INHIBITORS OF POLY(ADP-RIBOSE) POLYMERASES



(I)

(57) Abstract

Compounds of formula (I) are poly(ADP-ribosyl)transferase (PARP) inhibitors, and are useful as therapeutics in treatment of cancers and the amelioration of the effects of stroke, head trauma, and neurodegenerative disease. As cancer therapeutics, the compounds of the invention may be used, e.g., in combination with cytotoxic agents and/or radiation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TRICYCLIC INHIBITORS OF POLY(ADP-RIBOSE) POLYMERASES

This application claims priority from Provisional Application No. 60/115,431 filed January 11, 1999, which is incorporated herein by reference in its entirety.

5

FIELD OF THE INVENTION

The invention pertains to compounds that inhibit poly(ADP-ribose) polymerases, thereby retarding the repair of damage to DNA strands, and to methods of preparing such compounds. The invention also relates the use of such compounds in pharmaceutical compositions and therapeutic treatments useful for potentiation of anti-cancer therapies and inhibition of neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases.

10

BACKGROUND OF THE INVENTION

Poly(ADP-ribose) polymerases (PARPs), nuclear enzymes found in almost all eukaryotic cells, catalyze the transfer of ADP-ribose units from nicotinamide adenine dinucleotide (NAD⁺) to nuclear acceptor proteins, and are responsible for the formation of protein-bound linear and branched homo-ADP-ribose polymers. Activation of PARP and resultant formation of poly(ADP-ribose) can be induced by DNA strand breaks after exposure to chemotherapy, ionizing radiation, oxygen free radicals, or nitric oxide (NO).

15

20

Because this cellular ADP-ribose transfer process is associated with the repair of DNA strand breakage in response to DNA damage caused by radiotherapy or chemotherapy, it can contribute to the resistance that often develops to various types of cancer therapies. Consequently, inhibition of PARP may retard intracellular DNA repair and enhance the antitumor effects of cancer therapy. Indeed, *in vitro* and *in vivo* data show that many PARP inhibitors potentiate the effects of ionizing radiation or cytotoxic drugs such as DNA methylating agents. Therefore, inhibitors of the PARP enzyme are useful as cancer chemotherapeutics.

25

30

In addition, it has been shown that inhibition of PARP promotes resistance to brain injury after stroke (Endres et al., "Ischemic Brain Injury is Mediated by the

Activation of Poly(ADP-Ribose)Polymerase," *J. Cerebral Blood Flow Metab.* 17:1143-1151 (1997); Zhang, "PARP Inhibition Results in Substantial Neuroprotection in Cerebral Ischemia," *Cambridge Healthtech Institute's Conference on Acute Neuronal Injury: New Therapeutic Opportunities*, Sept. 18-24, 1998, Las Vegas, Nevada). The activation of PARP by DNA damage is believed to play a role in the cell death consequent to stroke, head trauma, and neurodegenerative diseases. DNA is damaged by excessive amounts of NO produced when the NO synthase enzyme is activated as a result of a series of events initiated by the release of the neurotransmitter glutamate from depolarized nerve terminals (Cosi et al., "Poly(ADP-Ribose) Polymerase Revisited: A New Role for an Old Enzyme: PARP Involvement in Neurodegeneration and PARP Inhibitors as Possible Neuroprotective Agents," *Ann. N.Y. Acad. Sci.*, 366-379). Cell death is believed to occur as a result of energy depletion as NAD⁺ is consumed by the enzyme-catalyzed PARP reaction. Therefore, inhibitors of the PARP enzyme are useful inhibitors of neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases.

Further, inhibition of PARP should be a useful approach for treatment of conditions or diseases associated with cellular senescence, such as skin aging, through the role of PARP in the signaling of DNA damage. See, e.g., U.S. Patent No. 5,589,483, which describes a method to extend the lifespan and proliferative capacity of cells comprising administering a therapeutically effective amount of a PARP inhibitor to the cells under conditions such that PARP activity is inhibited. Hence, inhibitors of the PARP enzyme are useful therapeutics for skin aging.

In yet a further application, PARP inhibition is being explored at the clinical level to prevent development of insulin-dependent diabetes mellitus in susceptible individuals (Saldeen et al., "Nicotinamide-induced apoptosis in insulin producing cells in associated with cleavage of poly(ADP-ribose) polymerase," *Mol. Cellular Endocrinol.* (1998), 139:99-107). PARP inhibitors should therefore be useful as diabetes-prevention therapeutics.

PARP inhibition is also an approach for treating inflammatory conditions such as arthritis (Szabo et al., "Protective effect of an inhibitor of poly(ADP-ribose)

synthetase in collagen-induced arthritis," *Portland Press Proc.* (1998), 15:280-281; Szabo, "Role of Poly(ADP-ribose) Synthetase in Inflammation," *Eur. J. Biochem.* (1998), 350(1):1-19; Szabo et al., "Protection Against Peroxynitrite-induced Fibroblast Injury and Arthritis Development by Inhibition of Poly(ADP-ribose) Synthetase," *Proc. Natl. Acad. Sci. USA* (1998), 95(7):3867-72). PARP inhibitors are therefore useful as therapeutics for inflammatory conditions.

Inhibition of PARP has usefulness for protection against myocardial ischemia and reperfusion injury (Zingarelli et al., "Protection against myocardial ischemia and reperfusion injury by 3-aminobenzamide, an inhibitor of poly (ADP-ribose) synthetase," *Cardiovascular Research* (1997), 36:205-215). Therefore, PARP inhibitors are useful in therapy of cardiovascular diseases.

The PARP family of enzymes is extensive. It has recently been shown that tankyrases, which bind to the telomeric protein TRF-1, a negative regulator of telomere length maintenance, have a catalytic domain that is strikingly homologous to PARP and have been shown to have PARP activity *in vitro*. It has been proposed that telomere function in human cells is regulated by poly(ADP-ribosyl)ation. PARP inhibitors have utility as tools to study this function. Further, as a consequence of regulation of telomerase activity by tankyrase, PARP inhibitors should have utility as agents for regulation of cell life-span, e.g., for use in cancer therapy to shorten the life-span of immortal tumor cells, or as anti-aging therapeutics, since telomere length is believed to be associated with cell senescence.

Competitive inhibitors of PARP are known. For example, Banasik et al. ("Specific Inhibitors of Poly(ADP-Ribose) Synthetase and Mono(ADP-Ribosyl)transferase," *J. Biol. Chem.* (1992) 267: 1569-1575) examined the PARP-inhibiting activity of 132 compounds, the most potent of which were 4-amino-1,8-naphthalimide, 6(5*H*)-phenanthridone, 2-nitro-6(5*H*)-phenanthridone, and 1,5-dihydroxyisoquinoline. Griffin et al. reported the PARP-inhibiting activity for a series of benzamide compounds (U.S. Patent No. 5,756,510; see also "Novel Potent Inhibitors of the DNA Repair Enzyme poly (ADP-ribose)polymerase (PARP)," *Anti-Cancer Drug Design* (1995), 10:507-514) and quinalozinone compounds

(International Publication No. WO 98/33802). Suto et al. reported PARP inhibition by a series of dihydroisoquinoline compounds ("Dihydroisoquinolines: The Design and Synthesis of a New Series of Potent Inhibitors of Poly(ADP-ribose) Polymerase," *Anti-Cancer Drug Design* (1991), 7:107-117). Griffin et al. have reported other


- 5 PARP inhibitors of the quinazoline class ("Resistance-Modifying Agents. 5. Synthesis and Biological Properties of Quinazoline Inhibitors of the DNA Repair Enzyme Poly(ADP-ribose) Polymerase (PARP)," *J. Med. Chem.*, ASAP Article 10.1021/jm980273t S0022-2623(98)00273-8; Web Release Date: December 1, 1998).

- 10 Nonetheless, there is still a need for small-molecule compounds that are potent PARP inhibitors, especially those that have physical and chemical properties desirable for pharmaceutical applications.

SUMMARY OF THE INVENTION

- 15 The present invention is directed to compounds that function as potent poly(ADP-ribosyl)transferase (PARP) inhibitors and are useful as therapeutics, especially in treatment of cancers and the amelioration of the effects of stroke, head trauma, and neurodegenerative disease. As cancer therapeutics, the compounds of the invention may be used in combination with DNA-damaging cytotoxic agents, for example, topotecan, irinotecan, or temozolomide, and/or radiation.

- 20 In particular, the present invention is directed to compounds of the general formula (I):



wherein:

- 25 R¹ is: H;
halogen;
cyano;

an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group (e.g., unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino, alkoxy, alkyl, and aryl groups
 5 unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, carboxy, and optionally substituted amino and ether groups (such as O-aryl)); or

-C(O)-R¹⁰, where R¹⁰ is: H; an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl
 10 group (e.g., unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and alkyl and aryl groups unsubstituted or substituted with one or more substituents selected from halo, hydroxy, nitro, and amino); or OR¹⁰⁰ or NR¹⁰⁰R¹¹⁰, where R¹⁰⁰ and R¹¹⁰ are each independently H or an optionally
 15 substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group (e.g., unsubstituted or substituted with one or more substituents selected from alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups unsubstituted or substituted with one or more substituents selected from halogen,
 20 hydroxy, nitro, amino, and alkyl and aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and optionally substituted amino groups);

R² is H or alkyl;

25 R³ is H or alkyl;

R⁴ is H, halogen or alkyl;

X is O or S;

Y is (CR⁵R⁶)(CR⁷R⁸)_n or N=C(R⁵), where:

n is 0 or 1;


R^5 and R^6 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group (e.g., unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and lower alkyl, lower alkoxy, or aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino); and

R^7 and R^8 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group (e.g., unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and lower alkyl, lower alkoxy, and aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino);

where when R^1 , R^4 , R^5 , R^6 , and R^7 are each H, R^8 is not unsubstituted phenyl.

The invention is also directed to pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates of such compounds. Preferred compounds of the formula (I) include those where R^2 and R^3 are each independently selected from H and methyl.

In a preferred embodiment, the inventive compounds include those of generic formula (II):



wherein:

p is 1 or 2;

R^{11} is H or alkyl;

R^{12} is halogen or an optionally substituted aryl, alkyl, alkenyl, alkynyl or acyl group $-C(O)-R^{10}$ as defined above;


R^{13} is H or alkyl; and

R^{14} is H or halogen;

- 5 as well as pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates of such compounds.

In preferred compounds of the formula (II), R^{11} and R^{13} are each independently selected from H and methyl. More preferably, the invention is directed to compounds of formula (II) where R^{11} and R^{13} are each H, and R^{12} is optionally substituted aryl, and to pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates of such compounds. In another preferred embodiment of compounds of formula (II), R^{11} and R^{13} are each H, and R^{12} is halogen or optionally substituted aryl.

15 In another preferred embodiment, the inventive compounds include those of generic formula (III) below, as well as pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates thereof:



wherein:

20 R^{15} is H, halogen, or an alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and alkyl and aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino;

25 R^{16} is H; halogen; cyano; or an alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and alkyl and aryl

groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino;

R^{17} is H or alkyl; and

R^{18} is H, halogen, or alkyl;

5 where R^{15} , R^{16} , R^{17} and R^{18} are not all H.

In preferred compounds of the formula (III), R^{15} is substituted phenyl or $(CH_2)_q$ aryl, where q is 1 or 2.

In other preferred compounds of the formula (III), R^{16} is substituted or unsubstituted aryl.

10 The present invention is also directed to a method of inhibiting PARP enzyme activity, comprising contacting the enzyme with an effective amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof. The compounds of the invention are potent PARP inhibitors and preferably have a PARP-inhibiting activity corresponding to a K_i of
15 100 μ M or less in the PARP enzyme inhibition assay.

The present invention is further directed to a method of potentiating the cytotoxicity of a cytotoxic drug or ionizing radiation, comprising contacting cells with an effective amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof, in
20 combination with a cytotoxic drug or ionizing radiation. The compounds of the invention preferably have a cytotoxicity potentiation activity corresponding to a PF_{50} of at least 1 in the cytotoxicity potentiation assay.

The present invention is also directed to pharmaceutical compositions comprising an effective PARP-inhibiting amount of a compound of formula (I), (II),
25 or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof, together with a pharmaceutically acceptable carrier therefor.

The invention also provides therapeutic interventions appropriate in disease or injury states where PARP activity is deleterious to a patient, the therapeutic methods comprising inhibiting PARP enzyme activity in the relevant tissue of the patient by
30 administering a compound of formula (I), (II), or (III), or a pharmaceutically

acceptable salt, prodrug, active metabolite, or solvate thereof. In one such therapeutic intervention method provided by the present invention, the effectiveness of a cytotoxic drug or radiotherapy administered to a mammal in the course of therapeutic treatment is improved by administering to the patient, e.g., a mammal in
5 need of treatment, an effective PARP-inhibiting amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof, in conjunction with the administration of the cytotoxic drug or radiotherapy.

Another therapeutic intervention method provided by the present invention is
10 for delaying the onset of cell senescence associated with skin aging in a human, comprising administering to fibroblast cells in the human an effective PARP-inhibiting amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof.


Yet another therapeutic intervention method provided by the present
15 invention is a method for reducing the neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases in a mammal by administering an effective amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof, to the mammal.

The compounds of the present invention provide a therapeutic approach to
20 treatment of inflammatory conditions, comprising administering an effective amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof, to a patient in need of treatment.

Yet a further therapeutic intervention method provided by the present invention is a cardiovascular therapeutic method for protecting against myocardial
25 ischemia and reperfusion injury in a mammal, comprising administering to the mammal an effective amount of a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof.


The present invention is further directed to methods of synthesizing the tricyclic compounds of formula (I), wherein a 4-carboalkoxy indole (IV) is converted
30 to an intermediate 3-substituted-4-carboalkoxy indole, thereby incorporating the


intended ring carbon atoms, terminally substituted with one nitrogen atom, usually in the form of a nitro group. Additional functional groups, such as formyl or acyl, may be incorporated at the 3-position in this step. The nitro group is reduced to an amine and cyclized upon the 4-carboalkoxy group in an amide-forming reaction to yield the tricyclic heterocycle. The synthetic methods may further comprise derivatization at N-1 and C-2. The 3-formyl or 3-acyl intermediates can be converted to nitrogen-containing intermediates or to tricyclic indoles with N-N bonds, such as the compounds of formula (III).



DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

15 PARP-Inhibiting Agents:

In accordance with a convention used in the art, the symbol  is used in structural formulas herein to depict the bond that is the point of attachment of the moiety or substituent to the core or backbone structure. In accordance with another convention, in some structural formulae herein the carbon atoms and their bound

20 hydrogen atoms are not explicitly depicted, e.g.,  represents a methyl group,

 represents an ethyl group,  represents a cyclopentyl group, etc.

As used herein, the term "alkyl" means a branched- or straight-chained (linear) paraffinic hydrocarbon group (saturated aliphatic group) having from 1 to 10 carbon atoms in its chain, which may be generally represented by the formula

25 C_kH_{2k+1} , where k is an integer of from 1 to 10. Examples of alkyl groups include

methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, pentyl, n-pentyl, isopentyl, neopentyl, and hexyl, and the simple aliphatic isomers thereof. A "lower alkyl" is intended to mean an alkyl group having from 1 to 4 carbon atoms in its chain.

5 The term "alkenyl" means a branched- or straight-chained olefinic hydrocarbon group (unsaturated aliphatic group having one or more double bonds) containing 2 to 10 carbons in its chain. Exemplary alkenyls include ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, isobutenyl, and the various isomeric pentenyls and hexenyls (including both cis and trans isomers).

10 The term "alkynyl" means a branched or straight-chained hydrocarbon group having one or more carbon-carbon triple bonds, and having from 2 to 10 carbon atoms in its chain. Exemplary alkynyls include ethynyl, propynyl, 1-butylnyl, 2-butylnyl, and 1-methyl-2-butylnyl.

15 The term "carbocycle" refers to a saturated, partially saturated, unsaturated, or aromatic, monocyclic or fused or non-fused polycyclic, ring structure having only carbon ring atoms (no heteroatoms, i.e., non-carbon ring atoms). Exemplary carbocycles include cycloalkyl, aryl, and cycloalkyl-aryl groups.

20 The term "heterocycle" refers to a saturated, partially saturated, unsaturated, or aromatic, monocyclic or fused or non-fused polycyclic, ring structure having one or more heteroatoms selected from N, O, and S. Exemplary heterocycles include heterocycloalkyl, heteroaryl, and heterocycloalkyl-heteroaryl groups.

25 A "cycloalkyl group" is intended to mean a non-aromatic monovalent, monocyclic or fused polycyclic, ring structure having a total of from 3 to 18 carbon ring atoms (but no heteroatoms). Exemplary cycloalkyls include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptyl, adamantyl, phenanthrenyl, and like groups.

A "heterocycloalkyl group" is intended to mean a non-aromatic monovalent, monocyclic or fused polycyclic, ring structure having a total of from 3 to 18 ring atoms, including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur.

Illustrative examples of heterocycloalkyl groups include pyrrolidinyl, tetrahydrofuryl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, aziridinyl, and like groups.

The term "aryl" means an aromatic monocyclic or fused polycyclic ring structure having a total of from 4 to 18, preferably 6 to 18, ring carbon atoms (no
5 heteroatoms). Exemplary aryl groups include phenyl, naphthyl, anthracenyl, and the like.

A "heteroaryl group" is intended to mean an aromatic monovalent, monocyclic or fused polycyclic, ring structure having from 4 to 18, preferably 5 to 18, ring atoms, including from 1 to 5 heteroatoms selected from nitrogen, oxygen,
10 and sulfur. Illustrative examples of heteroaryl groups include pyrrolyl, thienyl, oxazolyl, pyrazolyl, thiazolyl, furyl, pyridinyl, pyrazinyl, triazolyl, tetrazolyl, indolyl, quinolinyl, quinoxalinyl, and the like.

The term "optionally substituted" is intended to indicate that the specified group is unsubstituted or substituted by one or more suitable substituents, unless the
15 optional substituents are expressly specified, in which case the term indicates that the group is unsubstituted or substituted with the specified substituents. Unless indicated otherwise (e.g., by indicating that a specified group is unsubstituted), the various groups defined above may be generally unsubstituted or substituted (i.e., they are optionally substituted) with one or more suitable substituents.

20 The term "substituent" or "suitable substituent" is intended to mean any substituent for a group that may be recognized or readily selected by the artisan, such as through routine testing, as being pharmaceutically suitable. Illustrative examples of suitable substituents include hydroxy, halogen (F, Cl, I, or Br), oxo, alkyl, acyl, sulfonyl, mercapto, nitro, alkylthio, alkoxy, cycloalkyl, heterocycloalkyl, aryl,
25 heteroaryl, carboxy, amino (primary, secondary, or tertiary), carbamoyl, aryloxy, heteroaryloxy, arylthio, heteroarylthio, and the like (e.g., as illustrated by the exemplary compounds described herein). Suitable substituents are seen from the exemplary compounds that follow.

Preferred optional substituents for alkyl and aryl groups in the compounds of
30 the invention include halogens and aryl groups. Especially preferred for substituted

alkyl groups are perfluoro-substituted alkyls. Especially preferred optional substituents for aryl moieties include halogen, lower alkyl, -OH, -NO₂, -CN, -CO₂H, O-lower alkyl, aryl, -O-aryl, aryl-lower alkyl, -CO₂CH₃, -CONH₂, -OCH₂CONH₂, -NH₂, -SO₂NH₂, -OCHF₂, -CF₃, -OCF₃, and the like. Aryl moieties may also be
5 optionally substituted by two substituents forming a bridge, for example -O-(CH₂)_z-O-, where z is an integer of 1, 2, or 3.

A "prodrug" is intended to mean a compound that is converted under physiological conditions or by solvolysis, or metabolically, to a specified compound that is pharmaceutically active.

10 An "active metabolite" is intended to mean a pharmacologically active product produced through metabolism in the body of a specified compound.

A "solvate" is intended to mean a pharmaceutically acceptable solvate form of a specified compound that retains the biological effectiveness of such compound. Examples of solvates include compounds of the invention in combination with water,
15 isopropanol, ethanol, methanol, DMSO, ethyl acetate, acetic acid, or ethanolamine.

A "pharmaceutically acceptable salt" is intended to mean a salt that retains the biological effectiveness of the free-acid or base form of the specified compound and that is pharmaceutically suitable. Examples of pharmaceutically acceptable salts include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates,
20 monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates,
25 hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates, glycollates, tartrates, methane-sulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, and mandelates.

If an inventive compound is a base, a desired salt may be prepared by any
30 suitable method known in the art, including treatment of the free base with: an

inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with an organic acid, such as acetic acid, maleic acid, succinic acid, mandelic acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, pyranosidyl acid such as glucuronic acid or galacturonic acid; alpha-hydroxy acid such as citric acid or tartaric acid; amino acid such as aspartic acid or glutamic acid; aromatic acid such as benzoic acid or cinnamic acid; sulfonic acid such as p-toluenesulfonic acid or ethanesulfonic acid; or the like.

If an inventive compound is an acid, a desired salt may be prepared by any suitable method known in the art, including treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary, or tertiary), an alkali metal or alkaline earth metal hydroxide, or the like. Illustrative examples of suitable salts include: organic salts derived from amino acids such as glycine and arginine; ammonia; primary, secondary, and tertiary amines; and cyclic amines, such as piperidine, morpholine, and piperazine; as well as inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum, and lithium.


In the case of compounds, salts, or solvates that are solids, it is understood by those skilled in the art that the inventive compounds, salts, and solvates may exist in different crystalline or polymorph forms, all of which are intended to be within the scope of the present invention and specified formulas.

In some cases, the inventive compounds will have chiral centers. When chiral centers are present, the inventive compounds may exist as single stereoisomers, racemates, and/or mixtures of enantiomers and/or diastereomers. All such single stereoisomers, racemates, and mixtures thereof are intended to be within the broad scope of the generic structural formulae (unless otherwise indicated). Preferably, however, the inventive compounds are used in essentially optically pure form (as generally understood by those skilled in the art, an optically pure compound is one that is enantiomerically pure). Preferably, the compounds of the invention are at least 90% of the desired single isomer (80% enantiomeric excess), more preferably at least

95% (90% e.e.), even more preferably at least 97.5% (95% e.e.), and most preferably at least 99% (98% e.e.).



In some cases, compounds can occur in tautomeric forms. In such cases, it is intended that both tautomers are encompassed by the structural formulae.

- 5 The present invention is directed to the following PARP-inhibiting agents: compounds of the formula



- wherein R^1 , R^2 , R^3 , R^4 , X, and Y are as defined above; and pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates thereof. In preferred
10 embodiments, the PARP-inhibiting agents are compounds of the formula (I) where R^2 and R^3 are each independently H or methyl, and pharmaceutically acceptable salts, prodrugs, active metabolites, and solvates thereof.

More preferably, the agents are compounds of formula (II) or (III):



- 15 wherein the variables are as defined above, or pharmaceutically acceptable salts, prodrugs, active metabolites, or solvates thereof. In preferred embodiments for formula (II) and (III), R^{11} , R^{13} , and R^{17} are each independently H or methyl.

- 20 In a preferred embodiment, the inventive agents are compounds of formula (II) and pharmaceutically acceptable salts, prodrugs, active metabolites and solvates,


where R^{11} and R^{13} are each H, and R^{12} is an optionally substituted aryl group. In another preferred embodiment, the inventive agents are compounds of formula (III) and pharmaceutically acceptable salts, prodrugs, active metabolites and solvates, where R^{17} is H or methyl and R^{15} is optionally substituted aryl or alkyl.










5 In other preferred embodiments, R^{16} is substituted or unsubstituted aryl and R^{15} is hydrogen.

In other preferred embodiments, R^{16} is H, and R^{15} is substituted or unsubstituted aryl or alkyl.










Preferred compounds of the invention include:

10

















5









5
















5









10











5













5








5














5









10
















10



5








Pharmaceutical Methods and Compositions:

The invention is also directed to a method of inhibiting PARP enzyme activity, comprising contacting the enzyme with an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof. For example, PARP activity may be inhibited in mammalian tissue by administering a compound of formula (I) or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof. In addition to the compounds specified above, the following known compounds have been found to be useful for inhibiting PARP enzyme activity:



"Treating" or "treatment" is intended to mean mitigating or alleviating an injury or a disease condition in a mammal, such as a human, that is mediated by the inhibition of PARP activity, such as by potentiation of anti-cancer therapies or inhibition of neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases. Types of treatment include: (a) as a prophylactic use in a mammal, particularly when the mammal is found to be predisposed to having the disease condition but not yet diagnosed as having it; (b) inhibition of the disease condition; and/or (c) alleviation, in whole or in part, of the disease condition.

One treatment method involves improving the effectiveness of a cytotoxic drug or radiotherapy administered to a mammal in the course of therapeutic

treatment, comprising administering to the mammal an effective amount of an agent (compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate) in conjunction with administration of the cytotoxic drug (e.g., topotecan or irinotecan) or radiotherapy. The PARP-inhibiting agents may also be advantageously
5 used in a method for reducing neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases in a mammal by administering a therapeutically effective amount of an inventive agent to the mammal. The PARP-inhibiting agents of the invention may also be used in a method for delaying the onset of cell senescence associated with skin aging in a human, comprising administering to fibroblast cells in
10 the human an effective PARP-inhibiting amount of an agent. Further, the agents may also be used in a method for helping prevent the development of insulin-dependent diabetes mellitus in a susceptible individual, comprising administering a therapeutically effective amount of an agent. Additionally, the agents may also be employed in a method for treating an inflammatory condition in a mammal,
15 comprising administering a therapeutically effective amount of an agent to the mammal. Moreover, the agents may also be used in a method for treating cardiovascular disease in a mammal, comprising administering to the mammal a therapeutically effective amount of a PARP-inhibiting agent. As knowledge regarding the therapeutic roles of PARP inhibitors progresses in the art, other utilities
20 of the PARP-inhibiting agents of the invention will become apparent.

The activity of the inventive compounds as inhibitors of PARP activity may be measured by any of the suitable methods known or available in the art, including by *in vivo* and *in vitro* assays. An example of a suitable assay for activity measurements is the PARP enzyme inhibition assay described herein.

25 Administration of the compounds of the formula (I) and their pharmaceutically acceptable prodrugs, salts, active metabolites, and solvates may be performed according to any of the accepted modes of administration available in the art. Illustrative examples of suitable modes of administration include oral, nasal, parenteral, topical, transdermal, and rectal delivery. Oral and intravenous delivery
30 are preferred.

An inventive compound of formula (I) or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof may be administered as a pharmaceutical composition in any pharmaceutical form recognizable to the skilled artisan as being suitable. Suitable pharmaceutical forms include solid, semisolid, liquid, or lyophilized formulations, such as tablets, powders, capsules, suppositories, suspensions, liposomes, and aerosols. Pharmaceutical compositions of the invention may also include suitable excipients, diluents, vehicles, and carriers, as well as other pharmaceutically active agents (including other PARP-inhibiting agents), depending upon the intended use.

Acceptable methods of preparing suitable pharmaceutical forms of the pharmaceutical compositions are known or may be routinely determined by those skilled in the art. For example, pharmaceutical preparations may be prepared following conventional techniques of the pharmaceutical chemist involving steps such as mixing, granulating, and compressing when necessary for tablet forms, or mixing, filling, and dissolving the ingredients as appropriate to give the desired products for oral, parenteral, topical, intravaginal, intranasal, intrabronchial, intraocular, intraaural, and/or rectal administration.

Solid or liquid pharmaceutically acceptable carriers, diluents, vehicles, or excipients may be employed in the pharmaceutical compositions. Illustrative solid carriers include starch, lactose, calcium sulphate dihydrate, terra alba, sucrose, talc, gelatin, pectin, acacia, magnesium stearate, and stearic acid. Illustrative liquid carriers include syrup, peanut oil, olive oil, saline solution, and water. The carrier or diluent may include a suitable prolonged-release material, such as glyceryl monostearate or glyceryl distearate, alone or with a wax. When a liquid carrier is used, the preparation may be in the form of a syrup, elixir, emulsion, soft gelatin capsule, sterile injectable liquid (e.g., solution), or a nonaqueous or aqueous liquid suspension.

A dose of the pharmaceutical composition contains at least a therapeutically effective amount of a PARP-inhibiting agent (i.e., a compound of formula (I), (II), or (III), or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate

thereof), and preferably contains one or more pharmaceutical dosage units. The selected dose may be administered to a mammal, for example, a human patient, in need of treatment of a condition mediated by inhibition of PARP activity, by any known or suitable method of administering the dose, including: topically, for
5 example, as an ointment or cream; orally; rectally, for example, as a suppository; parenterally by injection; or continuously by intravaginal, intranasal, intrabronchial, intraaural, or intraocular infusion. A "therapeutically effective amount" is intended to mean the amount of an agent that, when administered to a mammal in need thereof, is sufficient to effect treatment for injury or disease condition mediated by inhibition of
10 PARP activity, such as for potentiation of anti-cancer therapies and inhibition of neurotoxicity consequent to stroke, head trauma, and neurodegenerative diseases. The amount of a given compound of the invention that will be therapeutically effective will vary depending upon factors such as the particular compound, the disease condition and the severity thereof, the identity of the mammal in need
15 thereof, which amount may be routinely determined by artisans.

It will be appreciated that the actual dosages of the PARP-inhibiting agents used in the pharmaceutical compositions of this invention will be selected according to the particular complex being used, the particular composition formulated, the mode of administration and the particular site, and the host and condition being
20 treated. Optimal dosages for a given set of conditions can be ascertained by those skilled in the art using conventional dosage-determination tests. For oral administration, e.g., a dose that may be employed is from about 0.001 to about 1000 mg/kg body weight, with courses of treatment repeated at appropriate intervals.

25 Synthetic Processes:

The present invention is further directed to methods of synthesizing the PARP-inhibiting agents by processes such as those set forth below for exemplary compounds of the invention. In the following examples, the structures of the compounds were confirmed by one or more of the following: proton magnetic
30 resonance spectroscopy, infrared spectroscopy, elemental microanalysis, mass


spectrometry, thin layer chromatography, high performance liquid chromatography, and melting point.

Proton magnetic resonance (^1H NMR) spectra were determined using a 300 megahertz Tech-Mag, Bruker Avance 300DPX, or Bruker Avance 500 DRX spectrometer operating at a field strength of 300 or 500 megahertz (MHz). Chemical shifts are reported in parts per million (ppm, δ) downfield from an internal tetramethylsilane standard. Alternatively, ^1H NMR spectra were referenced to residual protic solvent signals as follows: $\text{CHCl}_3 = 7.26$ ppm; DMSO = 2.49 ppm; $\text{C}_6\text{HD}_5 = 7.15$ ppm. Peak multiplicities are designated as follows: s = singlet; d = doublet; dd = doublet of doublets; t = triplet; q = quartet; br = broad resonance; and m = multiplet. Coupling constants are given in Hertz (Hz). Infrared absorption (IR) spectra were obtained using a Perkin-Elmer 1600 series or a Midac Corporation FTIR spectrometer. Elemental microanalyses were performed by Atlantic Microlab Inc. (Norcross, GA) or Galbraith Laboratories (Nashville, TN), and gave results for the elements stated within $\pm 0.4\%$ of the theoretical values. Flash column chromatography was performed using Silica gel 60 (Merck Art 9385). Analytical thin layer chromatography (TLC) was performed using precoated sheets of Silica 60 F254 (Merck Art 5719). Melting points (mp) were determined on a MelTemp apparatus and are uncorrected. All reactions were performed in septum-sealed flasks under a slight positive pressure of argon, unless otherwise noted. All commercial solvents were reagent-grade or better and used as supplied.

The following abbreviations may be used herein: Et_2O (diethyl ether); DMF (*N,N*-dimethylformamide); DMSO (dimethylsulfoxide); MeOH (methanol); EtOH (ethanol); EtOAc (ethyl acetate); THF (tetrahydrofuran); Ac (acetyl); Me (methyl); Et (ethyl); and Ph (phenyl).

The general reaction protocols described below may be used to prepare the compounds of the invention.

General Synthetic Scheme 1:




- In Scheme 1, 4-carbomethoxyindole A is formylated or acylated under various
- 5 Vilsmeier or Friedel-Crafts conditions to yield B, where R^{23} is CHO or COR^{24} . 4-Carbomethoxyindole A serves as substrate for a 1,4-addition reaction to yield the nitroethyl intermediate B, where R^{23} is $CHR^{25}CH_2NO_2$. Intermediate B, where R^{23} is CHO, is transformed to the corresponding oxime (R^{27} is $CH=NOH$) or nitroalkene

(R^{27} is $\text{CH}=\text{CHNO}_2$) **C**, which is then catalytically reduced to the aminoalkyl derivative **D**. Nitroethyl intermediate **B** is transformed directly to **D** (when R^{23} is $\text{CHR}^{25}\text{CH}_2\text{NO}_2$) by reduction in some cases. Compound **D** spontaneously cyclizes to tricyclic lactams **E** ($n = 2$) and **EE**. Exposure of intermediate **D** to basic conditions
 5 also leads to tricyclic lactams **E** and **EE**. Compound **E** is optionally N-alkylated to form N-alkylated **E** or halogenated to yield **F**. Intermediate **F** can be transformed via a metal-catalyzed reaction (typically with palladium as catalyst) into a number of different substituted tricyclic lactams **G**, where R^{29} is aryl, alkyl, alkenyl or alkynyl. **G** is optionally further modified at R^{22} , R^{29} and R^{30} .

10 Acyl-substituted compounds of formula **J** (e.g., compound 42) can be made by reaction with CO and the corresponding alcohol with Pd/C catalyst. The esters **J** may be further converted to other acyl derivatives by hydrolysis to the free acid, followed by activation to $-\text{C}(\text{O})-\text{Lv}$, where Lv is a leaving group, by standard
 15 methods (e.g., March, *Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 4th edition, August 1992, John Wiley & Sons, New York, ISBN 0471601802), and, for example, conversion to amides or other acyl derivatives by reactions generally known in the art. Alternatively, the esters **J** can be directly converted to amides by standard aminolysis reactions, e.g., by reaction with primary or secondary amines such as dimethylamine or pyrrolidine.

20 General Synthetic Scheme 2:




$R^{23} = \text{COR}^{24}$, ($R^{24} = \text{H, aryl, (CH)}_q\text{aryl}$), $q = 1 \text{ or } 2$ $R^{32} = \text{H, aryl, (CH}_2)_q\text{aryl}$

$R^{29} = \text{optionally substituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, or heteroaryl, or H.}$

In Scheme 2, intermediate **BB**, where R^{23} is CHO , $(\text{CO})\text{aryl}$, or $\text{CO}(\text{CH}_2)_q\text{aryl}$ where q is 1 or 2, is transformed to tricyclic acyl hydrazone **H** by reaction with hydrazine.

General Synthetic Scheme 3:



10

In Scheme 3, the **M**, where Lv includes, for example, I , Br , or triflate, is coupled with a substituted alkyne **T** using palladium and copper catalysts (See e.g. Sonogashira, K., Tohda, Y., Hagihara, N. *Tetrahedron Lett.* **1975**, *50*, 4467-4470, incorporated herein by reference). The intermediate **N** can be cyclized with palladium catalyst (See e.g. Arcadi, A., Cacchu, S., Marinellito, F. *Tetrahedron Lett.* **1989**, *30*, 2581-2584, incorporated herein by reference) to give **P** which is further modified as described in Scheme 1 to the intermediate **BB**.

15


EXAMPLES:

20

The invention is further described by reference to the following specific examples. Unless otherwise indicated, all percentages and parts are by weight, and all temperatures are in degrees Celsius.

Example A: 3,4-Dihydropyrrolo[4,3,2-de]isoquinolin-5-(1H)-one (1)

Compound 1 was prepared as described below according to the procedure of Demerson et al., *J. Med Chem.* (1974), 17:1140, starting from methyl indole-4-carboxylate.



(a) Methyl indole-4-carboxylate:

A solution of methyl 2-methyl-3-nitrobenzoate (9.85 g, 50.5 mmol) and dimethylformamide dimethyl acetal (20.1 mL, 151 mmol) in DMF (53 mL) was heated at 130°C for 8 hours (h). The solution was concentrated on a high-vacuum rotovap to give the benzoate enamine as a viscous dark-red oil, 12.2 g (97% yield).
¹H NMR (DMSO-*d*₆) δ 2.83 (s, 6H), 3.85 (s, 3H), 5.42 (d, 1H, *J* = 13.6 Hz), 6.41 (d, 1H, *J* = 13.6 Hz), 7.25 (t, 1H, *J* = 7.9 Hz), 7.76 (d, 1H, *J* = 7.9 Hz), 7.88 (d, 1H, *J* = 7.9 Hz).

A solution of the benzoate enamine (12.2 g, 48.4 mmol) in toluene (200 mL) was treated with 10% palladium-on-carbon (2.7 g), and the mixture was hydrogenated under 50 p.s.i. of hydrogen at room temperature for 1.5 h. The mixture was filtered through a pad of Celite, and the pad was rinsed with EtOAc. The crude product was purified by flash chromatography (3:1 hexanes:EtOAc) to yield methyl indole-4-carboxylate as a yellow solid, 6.89 g (81%). mp 68-70°C; ¹H NMR (DMSO-*d*₆) δ 3.95 (s, 3H), 7.02 (s, 1H), 7.25 (t, 1H, *J* = 7.6 Hz), 7.60 (s, 1H), 7.75 (d, 1H, *J* = 7.6 Hz), 7.80 (d, 1H, *J* = 7.6 Hz), 11.54 (bs, 1H).

20

(b) Intermediate J - methyl 3-formylindole-4-carboxylate:

A solution of methyl indole-4-carboxylate (250 mg, 1.43 mmol) in dichloroethane (2 mL) was treated with a solution of POCl₃-DMF (1.5 equivalent (eq)) at room temperature (rt). The orange solution was heated at 50°C for 1 hour.

5 The reaction solution was poured into ice-cold aqueous (aq.) NaOAc (1 g in 2 mL), the aqueous solution was adjusted to pH = 8 with 1M NaOH, and extracted with EtOAc (10 mL x 3). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated to give methyl 3-formyl-indole-4-carboxylate as an oil, 271 mg (93%). ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.68 (s, 3H), 7.16 (t, 1H, *J*

10 = 7.8 Hz), 7.40 (dd, 1H, *J* = 7.8, 0.8 Hz), 7.56 (d, 1H, *J* = 7.8, 0.8 Hz), 8.16 (d, 1H, *J* = 3.2 Hz), 10.00 (s, 1H), 12.30 (br s, 1H).

(c) Intermediate K - methyl 3-formylindole-4-carboxylate-oxime:

A mixture of J (2.5 g, 12.3 mmol), *N*-hydroxylamine hydrochloride (4.27 g, 61.4 mmol), NaOAc (5.04 g, 61.4 mmol), H₂O (25 mL), and MeOH (25 mL) was

15 stirred for 1 h at ~50°C. At this time the mixture was cooled to room temperature and concentrated under vacuum to remove the MeOH. Fifty mL of H₂O was added, and the solid was filtered and washed with additional H₂O. The pure white solid was dried under vacuum at 40°C (2.57 g, 95%). ¹H NMR (DMSO-*d*₆) δ 3.88 (s, 3H),

7.23 (t, 1H, *J* = 7.7 Hz), 7.59 (dd, 1H, *J* = 7.4, 1.1 Hz), 7.70 (dd, 1H, *J* = 8.1, 1.1 Hz),

20 8.01 (s, 1H), 8.52 (d, 1H, *J* = 3.0 Hz), 11.13 (s, 1H), 11.97 (bs, 1H).

(d) Intermediate L - methyl 3-aminomethylindole-4-carboxylate hydrochloride:

Dry HCl gas was added to a solution of oxime intermediate K (2.4 g, 11 mmol) in 130 mL MeOH. Under an argon atmosphere, 0.2 g of 10% Pd/C was

25 added. Using a three-way valve, the system was evacuated under vacuum. Hydrogen gas was introduced via a balloon, and the reaction mixture was vigorously stirred for 4 h. At this time the balloon was removed, and argon was reintroduced. The mixture was filtered and concentrated to give a solid which became violet in color. The solids were washed with Et₂O, protected from air and light, and placed under vacuum at


30 room temperature. The violet solid (2.5 g, 96%) was used without further

purification. ^1H NMR ($\text{DMSO}-d_6$) δ 3.89 (s, 3H), 4.31 (m, 2H), 7.23 (t, 1H, $J = 7.7$ Hz), 7.68 (d, 1H, $J = 2.6$ Hz), 7.74 (dd, 1H, $J = 8.1, 1.1$ Hz), 7.78 (dd, 1H, $J = 7.2, 1.1$ Hz), 8.05 (bs, 3H), 11.92 (bs, 1H).


(e) Compound 1 - 3,4-dihydropyrrolo[4,3,2-*de*]isoquinolin-5-(1*H*)-one:

5 A solution of intermediate L (2.4 g, 10.0 mmol) in 24 mL absolute EtOH was added to a methanolic solution of NaOMe (0.45 g Na, 24 mL anhydrous MeOH). After stirring at room temperature for 1.5 h, the mixture was concentrated under vacuum to give a residue. With stirring, ice-cold H_2O (75 mL) was added to the residue, and the solids were filtered and washed with cold H_2O (50 mL). Drying in a vacuum oven at 40°C afforded 1.51 g (87%) of analytically pure 1 as a tan solid. ^1H NMR ($\text{DMSO}-d_6$) δ 4.78 (s, 2H), 7.14 (t, 1H, $J = 7.7$ Hz), 7.18 (s, 1H), 7.30 (d, 1H, $J = 7.0$ Hz), 7.44 (d, 1H, $J = 8.1$ Hz), 7.59 (s, 1H), 11.13 (bs, 1H); HRMS ($\text{M}+\text{H}$), 173.0718; Anal. ($\text{C}_{10}\text{H}_8\text{N}_2\text{O} \cdot 0.2 \text{H}_2\text{O}$) C, H, N.


15 Example B: 2-Bromo-3,4-dihydropyrrolo[4,3,2-*de*]isoquinolin-5-(1*H*)-one (2)



A suspension of Compound 1 (0.086 g, 0.5 mmol) in 40 mL CH_2Cl_2 was treated with 90% pyridinium tribromide (0.267 g, 0.75 mmol) at 0°C . The reaction mixture was stirred at 0°C for 30 minutes (min.). The solvent was removed *in vacuo*, and ice-water was added to the residue. The resulting suspension was stirred
 20 vigorously at 0°C for 30 min. and then filtered, to give 0.068 g (54%) of a brown solid, which was used in the next step without further purification. IR (KBr) 3172, 1655, 1606, 1441, 1367, 1292, 755 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 4.61 (s, 2H), 7.17 (t, 1H, $J = 6.0$ Hz), 7.32 (d, 1H, $J = 6.0$ Hz), 7.39 (d, 1H, $J = 6.0$ Hz), 7.71 (s, 1H),
 25 11.92 (s, 1H); LRMS ($\text{M}+\text{H}$) 251/253.

Example C: Phenyl-3,4-dihydropyrrolo[4,3,2-de]isoquinolin-5-(1H)-one (3)

To a suspension of **2** (0.1065 g, 0.424 mmol) in 20 mL toluene/10 mL EtOH was added phenylboronic acid (0.08 g, 0.636 mmol), Na₂CO₃ (0.113 g, 1.06 mmol) dissolved in a minimum amount of water, LiCl (0.054 g, 1.27 mmol), and tetrakis(triphenylphosphine)palladium(0) (24.5 mg, 21.0 μmol). The reaction mixture was refluxed for 16 h. The solvent was removed *in vacuo*, and the residue was taken up in EtOAc and washed with saturated aqueous NaHCO₃, H₂O, and brine. The organic layer was dried over Na₂SO₄ and concentrated to give a yellow solid, which was purified by flash column chromatography eluting with a gradient of 20% of EtOAc in hexanes to give 0.098 g of a mixture of **3** as a yellow solid. mp 215-218°C (dec); ¹H NMR (DMSO-*d*₆) δ 5.04 (s, 2H), 7.17 (t, 1H, *J* = 7.5 Hz), 7.34 (d, 1H, *J* = 6.6 Hz), 7.35 (d, 1H, *J* = 7.4 Hz), 7.50 (m, 4H), 7.66 (d, 1H, *J* = 7.7 Hz), 7.84 (s, 1H), 11.64 (s, 1H); HRMS (M+H) 249.1023.


Example D: Compounds 4 and 5

To a suspension of Compound 2 in 30 mL toluene/15 mL EtOH was added 4-formylbenzeneboronic acid (0.457 g, 3.05 mmol), Na₂CO₃ (0.538 g, 5.08 mmol) dissolved in a minimum amount of water, LiCl (0.258 g, 6.09 mmol), and tetrakis(triphenylphosphine)palladium(0) (0.117 g, 0.102 mmol). The reaction mixture was refluxed for 48 h. The solvent was removed *in vacuo*, and the residue was taken up in EtOAc and washed with saturated aqueous NaHCO₃, H₂O, and brine. The organic layer was dried over MgSO₄ and concentrated to give a yellow solid, which was purified by flash column chromatography eluting with a gradient of 60-80% of EtOAc in CHCl₃ to give 0.370 g of a mixture of 4 and 5. Acetal 5 was converted to the aldehyde 4 using 5 mL MeOH/3 mL H₂O and a catalytic amount of conc. H₂SO₄.

4: IR (KBr) 1694, 1653, 1601, 1261, 821, 746 cm⁻¹; ¹H NMR (DMSO-*d*₆) δ 5.09 (s, 2H), 7.26 (t, 1H, *J* = 6.0 Hz), 7.36 (d, 1H, *J* = 6.0 Hz), 7.50 (d, 1H, *J* = 6.0 Hz), 7.85 (d, 2H, *J* = 9.0 Hz), 7.91 (s, 1H), 8.02 (d, 2H, *J* = 9.0 Hz), 10.01 (s, 1H), 11.86 (s, 1H); LRMS (M+H) 277.


5: ¹H NMR (DMSO-*d*₆) δ 1.15 (t, 6H, *J* = 6.0 Hz), 3.70 (q, 4H, *J* = 6.0 Hz), 5.03 (s, 2H), 5.51 (s, 1H), 7.20 (t, 1H, *J* = 6.0 Hz), 7.33 (d, 1H, *J* = 6.0 Hz), 7.46 (d, 1H, *J* = 6.0 Hz), 7.51 (d, 2H, *J* = 9.0 Hz), 7.65 (d, 2H, *J* = 9.0 Hz), 7.82 (s, 1H), 11.65 (s, 1H).

Example E: Compound 6



To a solution of 2M $(\text{CH}_3)_2\text{NH}$ in MeOH (0.81 mL, 1.61 mmol) was added 5N HCl-MeOH (0.11 mL, 0.536 mmol), followed by a suspension of the aldehyde 4 (0.074 g, 0.268 mmol) in 3 mL MeOH and NaBH_3CN (0.017 g, 0.268 mmol). The resulting suspension was stirred for 72 h at room temperature. Concentrated HCl was added until the pH was less than 2, and the MeOH was removed *in vacuo*. The residue was taken up in H_2O and extracted with EtOAc. The aqueous solution was brought to about pH 9 with solid KOH and extracted with EtOAc. The organic layer was dried over MgSO_4 and concentrated to give a yellow solid, which was purified by flash silica gel chromatography eluting with a gradient of 3% MeOH in CHCl_3 to 10% MeOH/ NH_3 in CHCl_3 , to give 0.023 g of an orange solid. ^1H NMR ($\text{DMSO}-d_6$) δ 2.17 (s, 6H), 3.44 (s, 2H), 5.04 (s, 2H), 7.19 (t, 1H, $J = 6.0$ Hz), 7.33 (d, 1H, $J = 6.0$ Hz), 7.42 (d, 1H, $J = 6.0$ Hz), 7.48 (d, 2H, $J = 9.0$ Hz), 7.63 (d, 2H, $J = 9.0$ Hz), 7.81 (s, 1H), 11.62 (s, 1H); LRMS ($\text{M}+\text{H}$) 306; Anal. ($\text{C}_{19}\text{H}_{19}\text{N}_3\text{O} \cdot 0.75 \text{H}_2\text{O}$) C, H, N.

15 Example F: Compounds 7 and 7a




Sixty percent sodium hydride (0.267 g, 6.67 mmol) was added to a solution of 1 (0.50 g, 2.9 mmol) in 7 mL DMF at 0°C . The reaction mixture was stirred at 0°C for 30 min., and then iodomethane (0.18 mL, 2.9 mmol) was added at 0°C . The reaction mixture was allowed to warm to room temperature and stirred for 1.5 h. The solvent was removed *in vacuo*, and the residue was taken in EtOAc and washed with H_2O and brine. The organic layer was dried over MgSO_4 and concentrated to give a brown solid, which was purified by flash silica gel chromatography eluting with a gradient of 0-1% of MeOH in CHCl_3 to give 0.270 g (50%) of 7 and 0.104 g (18%) of 7a, both as pale yellow solids.

7: IR (KBr) 3205, 1658, 1610, 1475, 1302, 1280, 817 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 3.80 (s, 3H), 4.76 (s, 2H), 7.15 (s, 1H), 7.18 (t, 1H, $J = 6.0$ Hz), 7.31 (d, 1H, $J = 6.0$ Hz), 7.51 (d, 1H, $J = 6.0$ Hz), 7.62 (s, 1H); LRMS ($\text{M}+\text{H}$) 187.

7a: IR (KBr) 1666, 1618, 1425, 1300, 1272, 1189, 742 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 3.05 (s, 3H), 3.81 (s, 3H), 4.89 (s, 2H), 7.17-7.22 (m, 2H), 7.35 (d, 1H, $J = 6.0$ Hz), 7.51 (d, 1H, $J = 6.0$ Hz); LRMS ($\text{M}+\text{H}$) 201.


Example G: Compound 9



10 Compound 9 was prepared from bromide 8 using a procedure similar to that described above for preparing Compound 4. IR (KBr) 1699, 1662, 1601, 1466, 1292, 1226 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 3.82 (s, 3H), 4.88 (s, 2H), 7.30 (t, 1H, $J = 6.0$ Hz), 7.39 (d, 1H, $J = 6.0$ Hz), 7.65 (d, 1H, $J = 6.0$ Hz), 7.78 (s, 1H), 7.82 (d, 2H, $J = 9.0$ Hz), 8.05 (d, 2H, $J = 9.0$ Hz), 10.08 (s, 1H); HRMS ($\text{M}+\text{H}$) 291.1130.


15

Example H: 3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (10)



Compound 10 was prepared according to a process generally described by Clark et. al (*J. Med. Chem.* (1990), 33:633-641) and Somei et. al (*Chem. Pharm. Bull.* (1988), 36:1162-1168).

20



Compound **M** was first prepared as follows. A solution of methyl indole-4-carboxylate (3.28 g, 18.7 mmol) and nitroethylacetate (2.99 g, 22.5 mmol) in xylenes (23 mL) was treated with 4-*t*-butylcatechol (22 mg) and heated at reflux for 3.5 h.

5 The solution was allowed to cool to room temperature and the solvent removed under reduced pressure. The residue was purified by flash chromatography (3:1

hexanes:EtOAc), to give a pale-yellow solid, 4.13 g (89%). mp 101-102°C; ¹H NMR (DMSO-*d*₆) δ 3.54 (t, 2H, *J* = 7.0 Hz), 3.93 (s, 3H), 4.79 (t, 2H, *J* = 7.0 Hz), 7.23 (m, 2H), 7.43 (s, 1H), 7.66 (m, 2H), 11.49 (bs, 1H); HRMS (*M*+*H*) Calcd for


10 C₁₂H₁₂N₂O₄+H: 249.0875, Found: 249.0870.

Intermediate **M** (1.12 g, 4.53 mmol) was dissolved in MeOH (70 mL) by gently heating. Aqueous 2M HCl (70 mL) was added. With vigorous stirring, 7.0 g of zinc dust was added portionwise, and the resulting mixture was heated at reflux for 30 min. The hot reaction mixture was filtered; the filtrate was treated with aqueous

15 2M NaOH (85 mL), and the resulting mixture was filtered through a paper-lined Buchner funnel. The filter cake was rinsed with MeOH. The MeOH was removed under reduced pressure, and the aqueous mixture was extracted with EtOAc (2 x 100 mL). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was crystallized with CH₂Cl₂/MeOH


20 to give the tricycle as a yellow solid, 611 mg (73%). mp 234-236°C; ¹H NMR (DMSO-*d*₆) δ 2.55 (m, 2H), 2.98 (m, 2H), 7.22 (t, 1H, *J* = 7.7 Hz), 7.31 (s, 1H), 7.58 (d, 1H, *J* = 7.7 Hz), 7.70 (d, 1H, *J* = 7.7 Hz), 8.04 (bt, 1H), 11.17 (bs, 1H); Anal. (C₁₁H₁₀N₂O) C, H, N.

Example I: 2-Bromo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (11)



Compound 10 (264 mg, 1.42 mmol) in CH_2Cl_2 (30 mL) and THF (30 mL) was treated with pyridinium tribromide (0.534 g, 1.67 mmol) at 0°C . The orange solution was stirred for 10 min., and then allowed to warm to ambient temperature and stirred for an additional hour. Water (30 mL) was added, and the organic solvents were removed *in vacuo*. The aqueous solution was adjusted to pH = 8-9 with 1M NaOH and extracted with CH_2Cl_2 (3 x 30 mL). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was recrystallized ($\text{CH}_2\text{Cl}_2/\text{MeOH}$) to yield the tricyclic bromide as a yellow solid, 305 mg (81%). mp $204\text{-}206^\circ\text{C}$ (dec); ^1H NMR ($\text{DMSO-}d_6$) δ 2.85 (m, 2H), 3.45 (m, 2H), 7.25 (t, 1H, $J = 7.8$ Hz), 7.52 (d, 1H, $J = 7.8$ Hz), 7.72 (d, 1H, $J = 7.8$ Hz), 8.14 (bt, 1H), 12.05 (bs, 1H); HRMS ($\text{M}+\text{H}$) Calcd for $\text{C}_{11}\text{H}_9\text{BrN}_2\text{O}+\text{H}$: 264.9976. Found: 264.9984.


Example J: 2-Phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (12)



Tricyclic bromide **11** (0.2 g, 0.75 mmol) in toluene (20 mL) and EtOH (10 mL) was treated with solid Na₂CO₃ (0.199 g, 1.88 mmol), LiCl (0.095 g, 2.25 mmol), 20 phenylboronic acid (0.138 g, 1.13 mmol), and water (0.50 mL). The solution was

degassed and tetrakis(triphenylphosphine)palladium(0) (43 mg, 5 mol %) was added. The solution was heated at reflux for 5 h, and then cooled to ambient temperature and diluted with water (20 mL). The aqueous layer was adjusted to pH = 7-8 with saturated aqueous K₂CO₃ and extracted with EtOAc (20 mL x 3). The organic solution was washed with water and brine, and dried (Na₂SO₄), filtered, and concentrated. The crude product was recrystallized (CH₂Cl₂/MeOH/hexanes) to yield the 2-phenyltricyclic as a pale-yellow solid, 183 mg (93%). mp 249-255°C (dec); ¹H NMR (CDCl₃/CD₄OD) δ 3.14 (m, 2H), 3.53 (m, 2H), 7.23 (t, 1H, *J* = 7.7 Hz), 7.33 (m, 1H), 7.44 (m, 2H), 7.55 (m, 3H), 7.83 (d, 1H, *J* = 7.7 Hz); HRMS (M+H) Calcd for C₁₇H₁₄N₂O+H: 263.1184, Found: 263.1189; Anal. (C₁₇H₁₄N₂O·0.8 H₂O) C, H, N.

Example K: 2-(4-Methoxyphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (13)



15


Tricyclic bromide 11 (48 mg, 0.18 mmol) in toluene (5 mL) and EtOH (2.5 mL) was treated with solid Na₂CO₃ (48 mg, 0.45 mmol), LiCl (23 mg, 0.54 mmol), *p*-methoxyphenylboronic acid (41 mg, 0.27 mmol), and water (0.25 mL). The solution was degassed, and tetrakis(triphenylphosphine)palladium(0) (10 mg, 5 mol %) was added. The solution was heated at reflux for 13 h, and then cooled to ambient temperature and diluted with water (10 mL). The aqueous layer was adjusted to pH = 7-8 with saturated aqueous K₂CO₃ and extracted with EtOAc (10 mL x 3). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was recrystallized (MeOH/THF) to yield the 2-(*p*-methoxyphenyl)tricyclic as a white solid, 47.4 mg (89%). mp 143-

25

148°C (dec); ^1H NMR (DMSO- d_6) δ 3.08 (m, 2H), 3.38 (m, 2H), 3.87 (s, 3H), 7.14 (d of ABq, 2H, $J = 8.6$ Hz), 7.22 (t, 1H, $J = 7.5$ Hz), 7.57 (d, 1H, $J = 7.5$ Hz), 7.64 (d of ABq, 2H, $J = 8.6$ Hz), 7.70 (d, 1H, $J = 7.5$ Hz), 8.11 (bt, 1H), 11.52 (bs, 1H); HRMS (M+H) Calcd for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2 + \text{H}$: 293.1290, Found: 293.1301; Anal.


5 (C₁₈H₁₆N₂O₂) C, H, N.

Example L: 2-(3-Nitrophenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4.3-cd]indol-6-one
(14)



10 Tricyclic bromide 11 (27 mg, 0.10 mmol) in 1,4-dioxane (1.0 mL) was treated with solid K_2CO_3 (41 mg, 0.30 mmol), *m*-nitrophenylboronic acid (34 mg, 0.20 mmol), and water (0.25 mL). The solution was degassed and tetrakis(triphenylphosphine)palladium(0) (12 mg, 10 mol %) was added. The solution was heated at 100°C for 1 h, then cooled to ambient temperature and diluted
15 with water (2 mL). The aqueous layer was adjusted to pH = 7-8 with saturated aqueous K_2CO_3 and extracted with EtOAc (5 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was purified by flash chromatography (3-5% MeOH in CHCl_3) to yield 14 as a yellow solid, 26.3 mg (87%). mp 268-270°C (dec.); ^1H NMR (DMSO- d_6) δ 3.16
20 (m, 2H), 3.45 (m, 2H), 7.33 (m, 1H), 7.65 (m, 1H), 7.76 (m, 1H), 7.78 (m, 1H), 8.30 (m, 1H), 8.53 (bs, 1H), 8.16 (m, 2H), 11.93 (bs, 1H); HRMS (M+Na) Calcd for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{O}_3 + \text{Na}$: 330.0855, Found: 330.0847; Anal. (C₁₇H₁₃N₃O₃·H₂O) C, H, N.


Example M: 2-(3-Hydroxymethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (16)



In a manner similar to that described above for Compound 12, the tricyclic
 5 bromide (381 mg, 1.44 mmol) and 3-formylbenzeneboronic acid (345 mg, 2.16
 mmol) were coupled to yield 2-(3-formylphenyl)-3,4,5,6-tetrahydro-1H-
 azepino[5,4,3-cd]indol-6-one **15**, 346 mg (83%), as a tan solid. ¹H NMR (300 MHz,
*d*₆-DMSO) δ 2.86 (m, 2H), 3.16 (m, 2H), 7.01 (t, 1H, *J* = 7.8 Hz), 7.34 (d, 1H, *J* =
 7.3 Hz), 7.50 (m, 2H), 7.73 (m, 2H), 7.85 (br t, 1H), 7.94 (s, 1H), 9.88 (s, 1H), 11.50
 10 (br s, 1H).


Compound **16** was isolated as a by-product from the reductive amination of
15 with dimethylamine and sodium cyanoborohydride, and recrystallized
 (CH₂Cl₂/hexanes), to give a pale-yellow solid. mp 258-259°C (dec); ¹H NMR
 (DMSO-*d*₆) δ 3.11 (m, 2H), 3.43 (m, 2H), 4.64 (d, 2H, *J* = 5.5 Hz), 5.36 (t, 1H, *J* =
 15 5.5 Hz), 7.26 (t, 1H, *J* = 7.6 Hz), 7.41 (m, 1H), 7.56 (m, 3H), 7.66 (m, 1H), 7.73 (d,
 1H, *J* = 7.6 Hz), 8.14 (m, 1H), 11.64 (bs, 1H); Anal. (C₁₈H₁₈N₂O₂·0.25 H₂O) C, H,
 N.

Example N: 2-(Phenylethynyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (17)



Tricyclic bromide 11 (58.6 mg, 0.22 mmol) in DMF (1 mL) was degassed and treated with tributyl(phenylethynyl)tin (95.2 mg, 0.24 mmol) and tetrakis(triphenylphosphine) palladium(0) (13 mg, 2 mol %). One crystal of 2,6-di-
5 butyl-4-methyl phenol was added, and the solution was heated at 60°C for 10 h. Starting material was still present, so the solution was heated at 100°C for an additional 2 h. The reaction mixture was cooled to ambient temperature and diluted with water (2 mL) and extracted with EtOAc (5 mL x 3). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude
10 product was purified by radial chromatography (2 mm SiO₂; 3% MeOH in CH₂Cl₂) to yield 17 as a white solid (34.8 mg, 55%). mp 255-256°C (dec); ¹H NMR (DMSO-*d*₆) δ 11.86 (s, 1H), 8.17 (m, 1H), 7.75 (d, 1H, *J* = 7.6 Hz), 7.63 (m, 3H), 7.51 (m, 3H), 7.33 (t, 1H, *J* = 7.6 Hz), 3.50 (m, 2H), 3.09 (m, 2H); HRMS (FAB, M+H) Calcd for C₁₉H₁₄N₂O+H: 287.1184, Found: 287.1192; Anal. (C₁₉H₁₄N₂O·0.6 H₂O) C, H,
15 N.


Example O: 1-Methyl-2-phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one
(18)



A solution of compound **12** (51.3 mg, 0.20 mmol) in THF (1 mL) and 0.1 mL 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (DMPU) was cooled with an ice/water bath and treated dropwise with a suspension of NaH (0.45 mmol) in THF (0.5 mL). The yellow mixture was allowed to stir at 0°C for 10 min., and was treated dropwise with a 1M solution of iodomethane in THF (0.22 mL, 0.22 mmol). The mixture was allowed to warm to ambient temperature and stirred for 30 min. The reaction was quenched at 0°C with saturated aqueous NH₄Cl, and extracted with EtOAc (5 mL x 3). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was purified by radial chromatography (2 mm SiO₂; 1-5% MeOH in CH₂Cl₂) to yield **18** as a white solid, 44.9 mg (81%). mp 254-256°C (dec.); ¹H NMR (DMSO-*d*₆) δ 2.88 (m, 2H), 3.40 (m, 2H), 3.74 (s, 3H), 7.34 (t, 1H, *J* = 7.7 Hz), 7.56 (m, 5H), 7.73 (d, 1H, *J* = 7.7 Hz), 7.80 (d, 1H, *J* = 7.7 Hz), 8.15 (bt, 1H); Anal. (C₁₈H₁₆N₂O·0.75 H₂O) C, H, N.

Compound **18a**, 1,5-dimethyl-2-phenyl-3,4,5,6-tetrahydro-1*H*-azepino[5,4,3-*cd*]indol-6-one, was isolated as a minor product. mp 175-177°C; ¹H NMR (DMSO-*d*₆) δ 2.91 (m, 2H), 3.19 (s, 3H), 3.65 (m, 2H), 3.75 (s, 3H), 7.34 (t, 2H, *J* = 7.8 Hz), 7.58 (m, 5H), 7.72 (d, 1H, *J* = 7.8 Hz), 7.79 (d, 1H, *J* = 7.8 Hz); Anal. (C₁₉H₁₈N₂O·0.5 H₂O) C, H, N.


Example P: 1-*N*-Methyl-3,4,5,6-tetrahydro-1*H*-azepino[5,4,3-*cd*]indol-6-one (**19**)



(19)

A solution of methyl indole-4-carboxylate (402 mg, 2.30 mmol) in DMF (5 mL) was cooled with an ice/water bath and treated with NaH (100 mg, 2.5 mmol, 60% in mineral oil). The resulting yellow solution was allowed to stir at 0°C for 30 min., then a solution of MeI (482 mg, 212 μ L, 3.4 mmol) in DMF (3.5 mL) was added dropwise. The solution was allowed to warm to ambient temperature. The reaction was quenched at 0°C with saturated aqueous NH_4Cl and extracted with EtOAc (10 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated to give methyl (N-methyl)-indole-4-carboxylate as a yellow oil, 430 mg (99%). The *N*-methyl carboxy indole was converted into the *N*-methyl-[5,6,7]-tricyclic indole in a manner similar to that described for Compound (10) to give 1-*N*-methyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one as a shiny white solid, 256 mg (54%, after recrystallization ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{hexanes}$)). mp 194-195°C; ^1H NMR (300 MHz, d_6 -DMSO) δ 2.96 (m, 2H), 3.43 (m, 2H), 3.82 (s, 3H), 7.29 (m, 2H), 7.64 (d, 1H, $J = 7.7$ Hz), 7.72 (d, 1H, $J = 7.7$ Hz), 8.09 (br t, 1H); HRMS (FAB, MH^+) Calcd for $\text{C}_{12}\text{H}_{13}\text{N}_2\text{O}$: 201.1028, Found: 201.1020; Anal. ($\text{C}_{12}\text{H}_{12}\text{N}_2\text{O} \cdot 0.2 \text{ H}_2\text{O}$) C, H, N.

Example Q: (*rac*)-3-Phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (20)




(20)

In a manner similar to that described for the preparation of methyl 3-(nitroethyl)-indole-4-carboxylate D above, methyl indole-4-carboxylate (85 mg, 0.49 mmol) and nitrostyrene (80 mg, 0.54 mmol) were heated at 160°C in a sealed tube for

12 h. The product was isolated by silica gel chromatography as a brown oil, 132 mg (83%). The intermediate nitro alkane was reduced/cyclized as described to give (*rac*)-3-phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-*cd*]indol-6-one as a white solid, 51.4 mg (48%, after chromatography and recrystallization). mp 201-203°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.73 (m, 2H), 4.42 (m, 1H), 7.28 (br m, 8H), 7.64 (d, 1H, *J* = 7.9 Hz), 7.77 (d, 1H, *J* = 7.9 Hz), 11.32 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₁₇H₁₅N₂O: 263.1184, Found: 263.1180; Anal. (C₁₇H₁₄N₂O·0.25 H₂O) C, H, N.


Example R: 2-(4-Fluorophenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-*cd*]indol-6-one
(23)



In a manner similar to that described for Compound 12, the tricyclic bromide (100 mg, 0.54 mmol) and 4-fluorobenzenboronic acid (79 mg, 0.57 mmol) were coupled to yield 2-(4-fluorophenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-*cd*]indol-6-one, 107 mg (99%), as a pale-yellow solid. ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.04 (m, 2H), 3.38 (m, 2H), 7.22 (app t, 1H, *J* = 7.7 Hz), 7.39 (m, 2H), 7.56 (dd, 1H, *J* = 8.0, 0.9 Hz), 7.64 (m, 3H), 8.05 (br t, 1H), 11.57 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₁₇H₁₄FN₂O: 281.1090, Found: 281.1093; Anal. (C₁₇H₁₃FN₂O·0.6 H₂O) C, H, N.


20

Example S: 8-Bromo-2-phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-*cd*]indol-6-one



A solution of Compound 12 (2-phenyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one) (22 mg, 0.08 mmol) in CH_2Cl_2 (1 mL) and THF (1 mL) was treated with pyridinium tribromide (29 mg, 0.09 mmol). The solution was stirred for 3 hours at room temperature and then diluted with water (2 mL), and the aqueous layer was
5 adjusted to pH = 9-10 with 1M NaOH. The mixture was extracted with CH_2Cl_2 (3 x 5 mL). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was purified by radial chromatography (1 mm silica gel; 1% MeOH in CHCl_3) to give the 8-bromo compound, 12.8 mg (47%), as a pale-yellow solid. ^1H NMR (300 MHz, d_6 -DMSO) δ 3.06 (m, 2H), 3.39 (m, 2H),
10 7.43 (app t, 1H, $J = 7.4$ Hz), 7.55 (app t, 2H, $J = 7.6$ Hz), 7.66 (app d, 2H, $J = 7.6$ Hz), 7.70 (app d, 1H, $J = 1.5$ Hz), 7.75 (app d, 1H, $J = 1.5$ Hz), 8.24 (br t, 1H), 11.77 (br s, 1H); HRMS (FAB, MH^+) Calcd for $\text{C}_{17}\text{H}_{14}\text{BrN}_2\text{O}$: 341.0289, Found: 341.0294.

15 Example T: 2-(4-(N,N-Dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (21)




In a manner similar to that described above for Compound 12, the tricyclic bromide (168 mg, 0.63 mmol) and 4-formylbenzeneboronic acid (142 mg, 0.95
20 mmol) were coupled to yield 2-(4-formylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 141 mg (77%), as a yellow solid. mp 238-240°C (dec.); ^1H NMR (300 MHz, d_6 -DMSO) δ 3.12 (m, 2H), 3.42 (m, 2H), 7.28 (t, 1H, $J = 7.6$ Hz), 7.59 (d, 1H, $J = 7.6$ Hz), 7.62 (d, 1H, $J = 7.6$ Hz), 7.88 (d of ABq, 2H, $J =$

7.7 Hz), 8.05 (d of ABq, 2H, $J = 7.7$ Hz), 8.11 (br t, 1H), 10.07 (s, 1H), 11.75 (br s, 1H); HRMS (FAB, MH^+) Calcd for $C_{18}H_{13}N_2O_2$: 291.1134, Found: 291.1132.

The aldehyde (310 mg, 1.07 mmol) in MeOH (40 mL) was treated with dimethyl amine (2M solution in MeOH, 6.41 mmol). The solution was cooled with an ice/water bath and treated dropwise with a solution of sodium cyanoborohydride (74 mg, 1.18 mmol) and zinc chloride (80 mg, 0.59 mmol) in MeOH (10 mL). The resulting solution was adjusted to pH = 6-7 with 2M methanolic HCl. After stirring for 30 min., the reaction was quenched with conc. HCl (0.2 mL) and the methanol was removed by evaporation. The residue was diluted with water (30 mL). The solution adjusted to pH = 10-11 with KOH (s) and extracted with CH_2Cl_2 (30 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was crystallized (CH_2Cl_2 /MeOH/hexanes) to give 2-(4-(N,N-dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 245 mg (72%), as an off-white solid. mp 226-229°C (dec.); 1H NMR (300 MHz, d_6 -DMSO) δ 2.18 (s, 6H), 3.06 (m, 2H), 3.40 (m, 2H), 3.44 (s, 2H), 7.21 (t, 1H, $J = 7.7$ Hz), 7.43 (d of ABq, 2H, $J = 7.9$ Hz), 7.56 (d, 1H, $J = 7.7$ Hz), 7.61 (d of ABq, 2H, $J = 7.9$ Hz), 7.69 (d, 1H, $J = 7.7$ Hz), 8.05 (br t, 1H), 11.53 (br s, 1H); HRMS (FAB, MH^+) Calcd for $C_{20}H_{22}N_3O$: 320.1763; Found: 320.1753; Anal. ($C_{20}H_{21}N_3O \cdot 0.55 H_2O$) C, H, N.

Example U: 2-(3-(N,N-Dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (22)




(22)

The aldehyde compound 15 (346 mg, 1.19 mmol) in MeOH (40 mL) was treated with dimethyl amine (2M solution in MeOH, 7.16 mmol). The solution was cooled with an ice/water bath and treated dropwise with a solution of sodium

cyanoborohydride (82 mg, 1.31 mmol) and zinc chloride (89 mg, 0.66 mmol) in MeOH (10 mL). The resulting solution was adjusted to pH = 6-7 with 2M methanolic HCl. After stirring for 30 min., the reaction was quenched with conc. HCl (0.2 mL) and the methanol was removed by evaporation. The residue was
5 diluted with water (30 mL). The solution was adjusted to pH = 10-11 with KOH (s) and extracted with CH₂Cl₂ (30 mL x 3). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was crystallized (CH₂Cl₂/MeOH/hexanes) to give 2-(3-(N,N-
dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one,
10 332 mg (87%), as shiny yellow crystals. mp 222-225°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.20 (s, 6H), 3.06 (m, 2H), 3.40 (m, 2H), 3.50 (s, 2H), 7.21 (t, 1H, *J* = 7.7 Hz), 7.41 (br d, 1H, *J* = 7.4 Hz), 7.50 (m, 4H), 7.69 (d, 1H, *J* = 7.1 Hz), 8.05 (br t, 1H), 11.56 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₂₀H₂₂N₃O: 320.1763, Found: 320.1753; Anal. (C₂₀H₂₁N₃O·0.25 H₂O) C, H, N.


15 Example V: Compound 25



To a solution of 2M (CH₃)₂NH in MeOH (0.6 mL, 1.13 mmol) was added 5N HCl-MeOH (0.08 mL, 0.380 mmol) followed by a suspension of the aldehyde (0.055 g, 0.188 mmol) in 3 mL MeOH and NaBH₃CN (0.012 g, 0.188 mmol). The resulting
20 suspension was stirred for 24 h at room temperature. Concentrated HCl was added until the pH was less than 2, and the MeOH was removed *in vacuo*. The residue was taken up in H₂O and extracted with EtOAc. The aqueous solution was brought to about pH 9 with solid (s) KOH and extracted with EtOAc. The organic layer was dried over MgSO₄ and concentrated to give a yellow solid, which was purified by
25 flash silica gel chromatography eluting with a gradient of CHCl₃ to 10% MeOH/NH₃ in CHCl₃ to give 0.024 g of a yellow solid. ¹H NMR (DMSO-*d*₆) 2.18 (s, 6H), 3.45

(s, 2H), 5.03 (s, 2H), 7.20-7.30 (m, 2H), 7.35 (d, 1H, $J = 6$ Hz), 7.40-7.58 (m, 3H), 7.60 (s, 1H), 7.79 (s br, 1H), 11.68 (s br, 1H); HRMS 306.1626.


Example W: 1,5-Dihydro-[1,2]diazepino[4,5,6-cd]-indol-6-one (27)



5

A solution of the intermediate J (3-formyl carboxy indole (246 mg, 1.21 mmol)) in MeOH (10 mL) and AcOH (0.1 mL) was treated with hydrazine hydrate (176 mg, 3.5 mmol) and the solution was heated at reflux for 30 min. The solution was cooled in an ice/water bath and the precipitated solid was collected by filtration to give 1,5-dihydro-[1,2]diazepino[4,5,6-cd]-indol-6-one, 168 mg (75%), as a bright-yellow solid. mp 335-336°C (dec.); ^1H NMR (300 MHz, d_6 -DMSO) δ 7.11 (t, 1H, $J = 7.8$ Hz), 7.44 (m, 3H), 7.56 (d, 1H, $J = 2.7$ Hz), 10.09 (s, 1H), 11.74 (br s, 1H); Anal. ($\text{C}_{10}\text{H}_7\text{N}_3\text{O}$) C, H, N.


15 Example X: 1,5-Dihydro-3-phenyl-[1,2]diazepino[4,5,6-cd]-indol-6-one (28)



A solution of methyl indole-4-carboxylate (40 mg, 0.23 mmol) in dichloroethane (2 mL) was treated with benzoyl chloride (0.69 mmol) at room temperature. The orange solution was cooled with an ice/water bath and treated with aluminum chloride (0.69 mmol). The dark-orange solution was warmed to room temperature over 1 hour, then poured into ice-cold aqueous 2M HCl. The aqueous solution was adjusted to pH = 9-10 with KOH (s), and extracted with CH_2Cl_2 (10 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was purified by radial chromatography (1 mm

silica gel; 3% MeOH in CHCl_3) to give methyl 3-phenacyl-indole-4-carboxylate as an oil, 63 mg (99%). A solution of the 3-phenacyl carboxy indole (60 mg, 0.25 mmol) in MeOH (5 mL) and conc. HCl (0.1 mL) was treated with hydrazine hydrate (36 mg, 0.73 mmol) and the solution was heated at reflux for 3 h. The reaction was quenched
5 with ice/water, and the aqueous layer was adjusted to pH = 10-11 with KOH (s) and extracted with CH_2Cl_2 (30 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated. The crude product was crystallized (CH_2Cl_2 /hexanes) to give 1,5-dihydro-3-phenyl-[1,2]diazepino[4,5,6-cd]-indol-6-one, 33 mg (51%), as a bright-yellow solid. mp 177-179°C; ^1H NMR (300 MHz, d_6 -
10 DMSO) δ 7.22 (m, 2H), 7.47 (m, 3H), 7.58 (m, 4H), 10.45 (s, 1H), 11.92 (br s, 1H); Anal. ($\text{C}_{10}\text{H}_7\text{N}_3\text{O} \cdot 0.75 \text{H}_2\text{O}$) C, H, N.

Example Y: 1,5-Dihydro-3-phenethyl-[1,2]diazepino[4,5,6-cd]-indol-6-one (29)




15 A solution of methyl indole-4-carboxylate (250 mg, 1.43 mmol) in dichloroethane (3 mL) was treated with 3-phenylpropionyl chloride (361 mg, 2.14 mmol) at room temperature. The orange solution was cooled to 0°C and treated with aluminum chloride (572 mg, 4.29 mmol). The reaction mixture was stirred at room temperature for 2 h, then poured into ice-cold 1M aqueous HCl. The aqueous
20 solution was adjusted to pH = 8 with 1M NaOH, and extracted with CH_2Cl_2 (10 mL x 3). The organic solution was washed with water and brine, dried (Na_2SO_4), filtered, and concentrated to give methyl 3-(3-phenylpropionyl)-indole-4-carboxylate as a pale-yellow solid, 395 mg (90%). A solution of the 3-(3-phenylpropionyl)-4-carboxy
25 indole (95.5 mg, 0.31 mmol) in MeOH (3 mL) and HCl (0.1 mL) was treated with hydrazine hydrate (47 mg, 0.93 mmol) and the solution was heated at reflux for 8 h. The solution was cooled in an ice/water bath and the precipitated solid was collected by filtration to give 1,5-dihydro-3-phenethyl-[1,2]diazepino[4,5,6-cd]-indol-6-one,

60.2 mg (71%). The crude product was purified by radial chromatography (2 mm SiO₂, 5:1 hexanes:EtOAc) to give a yellow solid. mp 182-183.5°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.80 (m, 2H), 2.84 (m, 2H), 7.22 (m, 2H), 7.31 (m, 4H), 7.54 (m, 2H), 7.81 (s, 1H), 10.19 (s, 1H), 11.92 (br s, 1H); Anal. (C₁₀H₇N₃O·0.1 H₂O) C, H, N.

5


Example Z: 2-(3-Trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (30)



In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 3-trifluoromethylphenylboronic acid (322 mg, 1.70 mmol) were coupled to yield 2-(3-trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 300 mg (80%), as a pale-yellow solid. mp 212.5-213.5°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.08 (m, 2H), 3.40 (m, 2H), 7.27 (app t, 1H, *J* = 7.8 Hz), 7.60 (d, 1H, *J* = 7.8 Hz), 7.71 (d, 1H, *J* = 7.5 Hz), 7.77 (m, 2H), 7.96 (m, 2H), 8.13 (br t, 1H), 11.78 (br s, 1H); MS (FAB, MH⁺) 331; Anal. (C₁₈H₁₃F₃N₂O·0.5 H₂O) C, H, N.

15

Example AA: 2-(4-Trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (31)




20

In a manner analogous to the method described above for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 4-trifluoromethylphenylboronic acid (322

mg, 1.70 mmol) were coupled to yield 2-(4-trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 261 mg (70%), as an off-white solid. mp 208-209°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.09 (m, 2H), 3.40 (m, 2H), 7.27 (app t, 1H, *J* = 7.8 Hz), 7.60 (dd, 1H, *J* = 8.1, 0.9 Hz), 7.71 (dd, 1H, *J* = 7.5, 0.6 Hz), 7.88 (m, 4H), 8.13 (br t, 1H), 11.77 (br s, 1H); MS (FAB, MH⁺) 331; Anal. (C₁₈H₁₃F₃N₂O·1.0 H₂O) C, H, N.

Example BB: 2-Benzofuran-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (32)



10


(32)

In a like manner to the example described above for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and benzo[b]furan-2-boronic acid (202 mg, 1.24 mmol) were coupled to yield 2-benzofuran-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 262 mg (77%), as a yellow solid. mp 207°C (dec.); ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.23 (m, 2H), 3.50 (m, 2H), 7.31 (m, 4H), 7.61 (dd, 1H, *J* = 8.1, 0.9 Hz), 7.70 (m, 3H), 8.14 (br t, 1H), 11.97 (br s, 1H); MS (FAB, MH⁺) 303; Anal. (C₁₉H₁₄N₂O₂·1.8 H₂O) C, H, N.

15

Example CC: 2-(3,5-bis-Trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (33)

20




(33)

In a manner similar to that described for preparation of Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 3,5-bis-trifluoromethylphenylboronic

acid (202 mg, 1.24 mmol) were coupled to yield 2-(3,5-bis-trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 70 mg (16%), as a pale-yellow solid. mp 230°C (dec.); ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.11 (m, 2H), 3.42 (m, 2H), 7.31 (app t, 1H, *J* = 7.8 Hz), 7.64 (d, 1H, *J* = 8.1 Hz), 7.73 (d, 1H, *J* = 7.5 Hz), 8.13 (br s, 1H), 8.16 (br t, 1H), 8.28 (br s, 2H), 11.95 (br s, 1H); MS (FAB, MH⁺) 399; Anal. (C₁₉H₁₂F₆N₂O·0.2 hexanes) C, H, N.

Example DD: 2-(4-Bromophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one
(34)




10

(34)

In a manner similar to that described for Compound 12, 2-iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (85 mg, 0.28 mmol; see Example NN below) and 4-bromophenylboronic acid (62 mg, 0.31 mmol) were coupled to yield 2-(4-bromophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 19 mg (20%), as a white solid. mp 160°C (dec.); ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.04 (m, 2H), 3.39 (m, 2H), 7.23 (app t, 1H, *J* = 7.5 Hz), 7.56 (dd, 1H, *J* = 8.1, 0.9 Hz), 7.60 (d, 2H, *J* = 8.7 Hz), 7.69 (dd, 1H, *J* = 7.5, 0.6 Hz), 7.73 (d, 2H, *J* = 8.4 Hz), 8.09 (br t, 1H), 11.64 (br s, 1H); MS (FAB, MH⁺) 341/343; Anal. (C₁₇H₁₃BrN₂O·0.6 H₂O) C, H, N.

15

Example EE: 2-(3-Chloro-4-fluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (35)




(35)

20

In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 3-chloro-4-fluorophenylboronic acid (217 mg, 1.24 mmol) were coupled to yield 2-(3-chloro, 4-fluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 217 mg (61%), as a pale-yellow solid. mp 234-235°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.04 (m, 2H), 3.39 (m, 2H), 7.24 (app t, 1H, *J* = 7.8 Hz), 7.57 (dd, 1H, *J* = 8.1, 0.9 Hz), 7.61 (m, 2H), 7.69 (dd, 1H, *J* = 7.5, 0.9 Hz), 7.85 (dd, 1H, *J* = 7.2, 2.1 Hz), 8.10 (br t, 1H), 11.68 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₁₇H₁₃ClFN₂O: 315.0700, Found: 315.0704; Anal. (C₁₇H₁₂ClFN₂O·1.0 H₂O·0.5 MeOH) C, H, N.

10

Example FF: 2-(4-*tert*-Butyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (36)




(36)

In a like manner as described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 4-*tert*-butylphenylboronic acid (302 mg, 1.70 mmol) were coupled to yield 2-(4-*tert*-butyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 150 mg (42%), as a white solid. mp 243-244°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 1.33 (s, 9H), 3.05 (m, 2H), 3.38 (m, 2H), 7.20 (app t, 1H, *J* = 7.8 Hz), 7.57 (m, 5H), 7.67 (dd, 1H, *J* = 7.2, 0.6 Hz), 8.07 (br t, 1H), 11.51 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₂₁H₂₃N₂O: 319.1810, Found: 319.1813; Anal. (C₂₁H₂₂N₂O·0.3 H₂O) C, H, N.


20

Example GG: 2-Phenyl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indole-6-thione (24)



Compound 12 (48.6 mg, 0.18 mmol) in toluene (2 mL) was treated with Lawesson's reagent (75 mg, 0.18 mmol) at room temperature. The solution was heated at reflux for 2 h, then allowed to cool to room temperature and diluted with water. The mixture was extracted with EtOAc (3 x 5 mL). The organic solution was washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. The crude product was crystallized (CH₂Cl₂/hexanes) to give the thioamide 34.4 mg (68%) as a yellow solid. mp 223-226°C (dec.); ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.10 (m, 2H), 3.50 (m, 2H), 7.23 (app t, 1H, *J* = 7.8 Hz), 7.57 (m, 1H), 7.61 (m, 3H), 7.69 (m, 2H), 8.19 (d, 1H, *J* = 7.6 Hz), 10.56 (br t, 1H), 11.68 (br s, 1H); HRMS (FAB, MH⁺) Calcd for C₁₇H₁₅N₂S: 279.0956, Found: 279.0952; Anal. (C₁₇H₁₄N₂S·0.25 H₂O) C, H, N, S.

Example HH: 2-Phenethyl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (37)




2-Phenylethynyl-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one (Compound 17) (37 mg, 0.13 mmol) and platinum oxide (1.5 mg, 0.05 mmol) were suspended in 2 mL MeOH under an argon atmosphere. The flask was flushed with hydrogen gas and the resulting mixture stirred at 24°C under 1 atmosphere of hydrogen for 20 h. The catalyst was filtered off and the resulting solution concentrated, leaving a pale-yellow crystalline solid. Purification by radial chromatography (5% MeOH in CHCl₃) followed by recrystallization (MeOH/CHCl₃/hexanes) yielded 2-phenethyl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 14 mg (37%), as a pale-yellow solid. mp 207-208°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.60 (m, 2H), 2.95 (m, 4H), 3.26 (m, 2H), 7.17 (m, 6H), 7.46 (dd,

1H, $J = 7.8, 0.6$ Hz), 7.61 (dd, 1H, $J = 7.5, 0.6$ Hz), 7.90 (br t, 1H), 11.16 (br s, 1H); MS (FAB, MH⁺) 291; Anal. (C₁₉H₁₈N₂O) C, H, N.


Example II: 2-(2-Chlorophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one

5 (38)



In a manner similar to that described for Compound 12, the tricyclic bromide (210 mg, 0.79 mmol) and 2-chlorophenylboronic acid (136 mg, 0.87 mmol) were coupled to yield 2-(2-chlorophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 78 mg (33%), as a shiny white solid. mp 275°C (dec); ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.76 (m, 2H), 3.38 (m, 2H), 7.23 (app t, 1H, $J = 7.8$ Hz), 7.56 (m, 5H), 7.71 (dd, 1H, $J = 7.5, 0.9$ Hz), 8.07 (br t, 1H), 11.53 (br s, 1H); MS (FAB, MH⁺) 297; Anal. (C₁₇H₁₃N₂OCl·0.15 H₂O) C, H, N.

15 Example JJ: 2-(2,4-Difluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (39)




In a manner similar to that described for Compound 12, the tricyclic bromide (200 mg, 0.75 mmol) and 2,4-difluorophenylboronic acid (131 mg, 0.83 mmol) were coupled to yield 2-(2,4-difluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 156 mg (69%), as a pale-yellow solid. mp 196-197°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.84 (m, 2H), 3.37 (m, 2H), 7.25 (app t, 1H, $J = 7.7$ Hz), 7.27 (m, 1H), 7.47 (m, 1H), 7.57 (dd, 1H, $J = 8.1, 0.9$ Hz), 7.64 (m, 1H), 7.70 (dd, 1H, $J = 7.5, 0.9$ Hz),

8.08 (br t, 1H), 11.58 (br s, 1H); MS (FAB, MH⁺) 299; Anal. (C₁₇H₁₂N₂OF₂·0.3 H₂O·0.37 CHCl₃) C, H, N.


Example KK: 2-(3-Chlorophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one

5 (40)



In a manner similar to that described for Compound 12, the tricyclic bromide (200 mg, 0.75 mmol) and 3-chlorophenylboronic acid (130 mg, 0.83 mmol) were coupled to yield 2-(3-chlorophenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 10 151 mg (67%), as a shiny pale-yellow solid. mp 147-149°C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.06 (m, 2H), 3.39 (m, 2H), 7.24 (app t, 1H, J = 7.8 Hz), 7.46 (m, 1H), 7.58 (m, 4H), 7.70 (m, 2H), 7.64 (m, 1H), 8.11 (br t, 1H), 11.68 (br s, 1H); MS (FAB, MH⁺) 297; Anal. (C₁₇H₁₃N₂OCl·0.9 H₂O) C, H, N.


15 Example LL: 2-Naphthalen-1-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one
(41)



In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 1-naphthaleneboronic acid (214 mg, 1.24 mmol) were 20 coupled to yield 2-naphthalen-1-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 70 mg (20%), as an off-white solid. mp 305°C (dec.); ¹H NMR (300 MHz, d₆-DMSO) δ 2.70 (m, 2H), 3.38 (m, 2H), 7.25 (app t, 1H, J = 7.5 Hz), 7.61 (m, 5H), 7.75


(dd, 1H, $J = 7.5, 0.9$ Hz), 7.82 (m, 1H), 8.06 (m, 3H), 11.67 (br s, 1H); MS (FAB, MH⁺) 313; Anal. (C₂₁H₁₆N₂O·0.2 H₂O) C, H, N.

Example MM: 6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methyl ester (42)






2-Iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (85 mg, 0.28 mmol; prepared as described below), palladium tetrakis(triphenylphosphine) (19 mg, 0.02 mmol), and triethylamine (52 mg, 0.51 mmol) were combined in toluene:methanol (8:2 (v/v), 2 mL). Carbon monoxide gas was bubbled through the mixture for 10 min. The reaction was then heated at 85°C in a sealed tube for 16 h. The solvent was evaporated and the orange solid purified by radial chromatography (chloroform to 5% methanol in chloroform). The white solid was recrystallized (chloroform/methanol/hexanes) to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methyl ester, 39 mg (100%), as an off-white solid. mp 266-267°C; ¹H NMR (300 MHz, *d*₆-DMSO) δ 3.25 (m, 2H), 3.43 (m, 2H), 3.89 (s, 3H), 7.38 (app t, 1H, $J = 7.8$ Hz), 7.61 (dd, 1H, $J = 8.1, 0.9$ Hz), 7.74 (dd, 1H, $J = 7.5, 0.9$ Hz), 8.17 (br t, 1H), 11.93 (br s, 1H); MS (FAB, MH⁺) 245; Anal. (C₁₃H₁₂N₂O₃) C, H, N.

Example NN: Preparation of 2-Iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (43)




1,3,4,5-Tetrahydro-azepino[5,4,3-cd]indol-6-one (620 mg, 3.35 mmol) was suspended in 80 mL THF/CH₂Cl₂ (1:1), and then cooled in an ice bath. Bis(trifluoroacetoxy)-iodo]benzene (1.73 g, 4.02 mmol) and iodine (850 mg, 3.35 mmol) were added and the reaction stirred at 0°C for 25 min. The ice bath was removed and the reaction allowed to stir for another 30 min. as it warmed to room temperature. The reaction was quenched by addition of aqueous sodium bisulfite. The layers were separated, and the organic layer was dried over MgSO₄, filtered, and concentrated *in vacuo* leaving a yellow solid. The crude solid was purified by flash chromatography (5% MeOH/CHCl₃) to yield 1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 308 mg (30%), as a pale-yellow solid: ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.79 (m, 2H), 3.40 (m, 2H), 7.14 (app t, 1H, *J* = 7.8 Hz), 7.46 (dd, 1H, *J* = 7.8, 0.6 Hz), 7.64 (dd, 1H, *J* = 7.5, 0.9 Hz), 8.06 (br t, 1H), 11.80 (br s, 1H); MS (FAB, MH⁺) 313.

By following methods analogous to those described in the above examples, the following compounds were also prepared:



Example OO:


2-(4-(*N*-Methylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one



- The *p*-aldehyde (150 mg, 0.52 mmol) prepared as described for compound 21 in MeOH (20 mL) was treated, as described, with methyl amine (8.03 M solution in EtOH, 3.10 mmol) and a solution of sodium cyanoborohydride (0.57 mmol) and zinc chloride (0.28 mmol) in MeOH (2 mL) to give, after recrystallization (isopropyl alcohol/hexanes), 2-(4-(*N*-methylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 108 mg (68%) as a yellow solid: m.p. 208-210°C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.34 (s, 3H), 3.05 (m, 2H), 3.39 (m, 2H), 3.77 (s, 2H), 7.20 (t, *J* = 7.7 Hz, 1H), 7.54 (m, 3H), 7.61 (d of ABq, *J* = 8.4 Hz, 2H), 7.67 (d, *J* = 7.6 Hz, 1H), 8.07 (br t, 1H), 11.55 (br s, 1H). HRMS (FAB, MH⁺) Calcd for C₁₉H₂₀N₃O: 306.1606. Found: 306.1601. Anal. (C₁₉H₁₉N₃O·0.4 H₂O) C, H, N.

15 Example PP:

2-(3-(*N*-Methylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one




In a manner similar to that described for Compound 22, the aldehyde 15 (200 mg, 0.69 mmol) in MeOH (20 mL) was treated with methyl amine (2.0 M solution in THF, 4.20 mmol) and a solution of sodium cyanoborohydride (0.76 mmol) and zinc chloride (0.38 mmol) in MeOH (1.4 mL) to give, after recrystallization

- 5 (CH₂Cl₂/MeOH/hexanes), 2-(3-(N-methylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 103 mg (49%) as pale yellow powder: m.p. 190-192 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.37 (s, 3H), 3.07 (m, 2H), 3.40 (m, 2H), 3.82 (s, 2H), 7.22 (t, *J* = 7.7 Hz, 1H), 7.39 (br d, 1H), 7.49 (m, 1H), 7.56 (m, 2H), 7.68 (m, 2H), 8.09 (br t, 1H), 11.61 (br s, 1H). HRMS (FAB, MH⁺) Calcd for
- 10 C₁₉H₂₀N₃O: 306.1606. Found: 306.1601. Anal. (C₁₉H₁₉N₃O·0.6 H₂O) C, H, N.


Example QQ:

1,5-Dihydro-3-methyl-[1,2]diazepino[4,5,6-cd]-indol-6-one




15

- In a manner similar to that described for compound 28, a solution of methyl indole-4-carboxylate (427 mg, 2.44 mmol) in dichloroethane (7 mL) was treated with acetyl chloride (0.5 mL) and aluminum chloride (130 mg). The intermediate ketone
- 20 (198 mg, 0.92 mmol) in MeOH (5 mL) and conc. HCl (0.05 mL) was treated, as described, with hydrazine hydrate (0.1 mL). The product precipitated, was collected by filtration and rinsed with ice-cold MeOH to give 1,5-dihydro-3-methyl-
- [1,2]diazepino[4,5,6-cd]-indol-6-one, 168 mg (92%) as a bright yellow solid: m.p. 335-336°C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.17 (s, 3H), 7.19 (t, *J* = 7.8 Hz, 1H),
- 25 7.54 (m, 2H), 7.67 (d, *J* = 2.8 Hz, 1H), 10.12 (s, 1H), 11.90 (br s, 1H). HRMS (FAB, MH⁺) Calcd for C₁₁H₁₀N₃O: 200.0824. Found: 200.0827. Anal. (C₁₁H₉N₃O) C, H, N.

Example RR:**2-(3-Aminophenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one**

5

In a manner similar to that described for Compound 12, the tricyclic bromide (428 mg, 1.61 mmol) and 3-aminobenzeneboronic acid monohydrate (300 mg, 1.94 mmol) were coupled to yield 2-(3-aminophenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 110 mg (25%) as an off-white solid: ¹H NMR (300 MHz, d₆-DMSO) δ 3.03 (m, 2H), 3.39 (m, 2H), 5.24 (s, 2H), 6.59 (br d, 1H), 6.78 (d, J = 7.7 Hz, 1H), 6.84 (m, 2H), 7.18 (m, 2H), 7.52 (d, J = 7.9 Hz, 1H), 7.66 (d, J = 7.4 Hz, 1H), 8.04 (br t, 1H), 11.41 (br s, 1H). HRMS (FAB, MH⁺) Calcd for C₁₇H₁₆N₃O: 278.1293. Found: 278.1297. Anal. (C₁₇H₁₅N₃O·1.1 H₂O) C, H, N.

15 Example SS:**2-(3-(3-Piperidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one**


20

In a manner similar to that described for Compound 22, the aldehyde 15 (109 mg, 0.38 mmol) in MeOH (10 mL) was treated with piperidine (0.19 mL, 1.9 mmol) and a solution of sodium cyanoborohydride (0.57 mmol) and zinc chloride (0.28 mmol) in MeOH (1.1 mL) to give, after recrystallization (CH₂Cl₂/hexanes), 2-(3-(3-piperidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 94.1 mg (69%) as pale yellow powder: m.p. 235-237 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 1.41 (m, 2H), 1.52 (m, 4H), 2.37 (m, 4H), 3.06 (m, 2H), 3.39 (m, 2H), 3.52 (s, 2H), 7.21 (t, *J* = 7.7 Hz, 1H), 7.31 (m, 1H), 7.54 (m, 4H), 7.69 (m, 1H), 8.08 (br t, 1H), 11.58 (br s, 1H). Anal. (C₂₃H₂₅N₃O·0.65 H₂O) C, H, N.

10

Example TT:

N-[3-(6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-phenyl]-acetamide




In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 3-acetamidophenylboronic acid (304 mg, 1.70 mmol) were coupled to yield N-[3-(6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-phenyl]-acetamide, 10 mg (3%) as a clear solid: m.p. 300.5-302.0 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.09 (s, 3H), 3.05 (m, 2H), 3.36 (m, 2H), 7.21 (app t, *J* = 7.8 Hz, 1H), 7.33 (d, *J* = 7.5 Hz, 1H), 7.44 (t, *J* = 7.8 Hz, 1H), 7.57 (m, 2H), 7.68 (d, *J* = 7.5 Hz, 1H), 7.92 (br s, 1H), 8.08 (br t, 1H), 10.10 (br s, 1H), 11.56 (br s, 1H). MS (FAB, MH⁺) 320. Anal. (C₁₉H₁₇N₃O₂) C, H, N.

20

Example UU:

2-[3-(4-Fluoro-phenoxy)-phenyl]-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one


25



In a manner similar to that described for Compound 12, the tricyclic bromide (200 mg, 0.75 mmol) and 3-(4-fluorophenoxy)-phenylboronic acid (213 mg, 0.83 mmol) were coupled to yield 2-[3-(4-fluorophenoxy)-phenyl]-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 170 mg (60%) as a yellow crystalline solid: m.p. 240-241 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.01 (m, 2H), 3.38 (m, 2H), 6.99 (m, 2H), 7.21 (m, 6H), 7.42 (m, 1H), 7.54 (m, 2H), 7.68 (m, 1H), 8.09 (br t, 1H), 11.60 (br s, 1H). MS (FAB, MH⁺) 373. Anal. (C₂₃H₁₇N₂O₂ F·0.5 H₂O) C, H, N.

10 Example VV:

2-Biphenyl-4-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one




In a manner similar to that described for Compound 12, the tricyclic bromide (150 mg, 0.57 mmol) and 2-Biphenyl-4-boronic acid (123 mg, 0.62 mmol) were coupled to yield 2-Biphenyl-4-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 87 mg (45%) as a pale yellow solid: m.p. 277-279 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.11 (m, 2H), 3.41 (m, 2H), 7.23 (app t, *J* = 7.8 Hz, 1H), 7.40 (m, 1H), 7.51 (app t, *J* = 7.2 Hz, 2H), 7.58 (d, *J* = 8.1 Hz, 1H), 7.77 (m, 7H), 8.10 (br t, 1H), 11.64 (br s, 1H). MS (FAB, MH⁺) 339 Anal. (C₂₃H₁₈N₂O·1.15 H₂O) C, H, N.

20


Example WW:

2-(4-Chloro-3-trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Compound 12, the tricyclic bromide (100 mg, 0.38 mmol) and 4-chloro-3-trifluoromethyl-phenylboronic acid (150 mg, 0.45 mmol) were coupled to yield 2-(4-chloro-3-trifluoromethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 121 mg (88%) as a pale yellow solid: m.p. 118.5-119 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.06 (m, 2H), 3.41 (m, 2H), 7.27 (app t, *J* = 7.8 Hz, 1H), 7.60 (dd, *J* = 7.8, 0.9 Hz, 1H), 7.73 (dd, *J* = 7.2, 0.9 Hz, 1H), 7.89 (m, 2H), 8.08 (d, *J* = 1.5 Hz, 1H), 8.14 (br t, 1H), 11.82 (br s, 1H). MS (FAB, MH⁺) 365. Anal. (C₁₈H₁₂ClF₃N₂O·0.45 H₂O·0.2 CHCl₃) C, H, N.


10

Example XX:**2-Naphthalen-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**

In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 2-naphthaleneboronic acid (214 mg, 1.24 mmol) were coupled to yield 2-Naphthalen-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 130 mg (37%) as a pale yellow solid: m.p. 261-262 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.18 (m, 2H), 3.42 (m, 2H), 7.24 (app t, *J* = 7.5 Hz, 1H), 7.58 (m, 3H), 7.72 (dd *J* = 7.5, 0.9 Hz, 1H), 7.84 (dd *J* = 8.4, 1.5 Hz, 1H), 8.07 (m, 5H), 11.74 (br s, 1H). MS (FAB, MH⁺) 313. Anal. (C₂₁H₁₆N₂O·0.9 H₂O) C, H, N.

20

Example YY:**2-[4-(2-Diethylamino-ethyl)-phenyl]-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one**



(As described in *Tet. Lett.* 1997 p. 3841) [2-(4-Bromo-phenyl)-ethyl]-diethyl-amine (256 mg, 1.00 mmol), diboron pinacol ester (279 mg, 1.10 mmol), 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium (24 mg, 0.03 mmol), and potassium acetate (294 mg, 3.00 mmol) were combined in a schlenk tube. The vessel was evacuated then refilled with argon thrice. Degassed DMF (6 mL) was added and the mixture stirred at 80 °C under an argon atmosphere for 2 h. 2-Bromo—3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one (239 mg, 0.90 mmol), a second portion of 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium (24 mg, 0.03 mmol), and sodium carbonate (2.5 mL of a 2.0 M aqueous solution, 5.00 mmol) were then added and the reaction stirred under an argon atmosphere at 80 °C for another 17 h. The reaction mixture was then poured into 25 mL water then extracted with 25 % IPA/CHCl₃ (3 x 20 mL). The combined organic extracts were dried (MgSO₄) and concentrated in vacuo leaving a brown oil. The crude product was passed through a short silica plug with 25 % MeOH/ CHCl₃ then purified by radial chromatography eluting with 20 % MeOH/ CHCl₃. Crystallization from MeOH/ CHCl₃/hexanes yielded 2-[4-(2-diethylamino-ethyl)-phenyl]-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one, 69 mg (19%) as a white solid: m.p. 224-224.5 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 0.98 (t, *J* = 6.9 Hz, 6H), 2.53 (q, *J* = 7.2 Hz, 4H), 2.69 (m, 4H), 3.04 (m, 2H), 3.37 (m, 2H), 7.19 (t, *J* = 7.8 Hz, 1H), 7.36 (d, *J* = 8.1 Hz, 2H), 7.55 (m, 3H), 7.88 (dd, *J* = 7.5, 0.9 Hz, 1H), 8.06 (br t, 1H), 11.51 (br s, 1H). MS (FAB, MH⁺): 362. Anal. (C₂₃H₂₇N₃O) C, H, N.


Example ZZ:

25 2-[3-(2-Hydroxy-ethyl)-phenyl]-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one

7.51 (dd, $J = 7.8, 0.9$ Hz, 1H), 7.70 (dd, $J = 7.5, 0.9$ Hz, 1H), 8.03 (br t, 1H), 11.39 (br s, 1H). MS (FAB, MH^+): 321. Anal. ($C_{20}H_{20}N_2O_2 \cdot 0.1 CH_2Cl_2$) C, H, N.

Example CCC:


5 **2-(4-Hydroxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**



In a manner similar to that described for Compound YY, 4-iodophenol (220 mg, 1.00 mmol), diboron pinacol ester (279 mg, 1.10 mmol), 1,1'-bis(diphenyl phosphino)ferrocenedichloro palladium (24 mg, 0.03 mmol), and potassium acetate
10 (294 mg, 3.00 mmol), 2-bromo-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one (239 mg, 0.90 mmol), a second portion of 1,1'-bis(diphenylphosphino)ferrocene dichloropalladium (24 mg, 0.03 mmol), and sodium carbonate (2.5 mL of a 2.0 M aqueous solution, 5.00 mmol) were reacted to yield 2-(4-hydroxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 39 mg (15%) as a beige solid: m.p. 300 °C
15 (dec); 1H NMR (300 MHz, d_6 -DMSO) δ 3.00 (m, 2H), 3.37 (m, 2H), 6.92 (d, $J = 8.7$ Hz, 2H), 7.16 (t, $J = 7.8$ Hz, 1H), 7.49 (m, 3H), 7.65 (dd, $J = 7.5, 0.9$ Hz, 1H), 8.04 (br t, 1H), 9.73 (br s, 1H), 11.40 (br s, 1H). MS (electrospray, MH^+): 279. Anal. ($C_{17}H_{14}N_2O_2$) C, H, N.

20 Example DDD:


2-(2-Hydroxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Example YY, 2-iodophenol (220 mg, 1.00 mmol), diboron pinacol ester (279 mg, 1.10 mmol), 1,1'-bis(diphenyl phosphino)ferrocenedichloro palladium (24 mg, 0.03 mmol), and potassium acetate (294 mg, 3.00 mmol), 2-bromo-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one (239 mg, 0.90 mmol), a second portion of 1,1'-bis(diphenylphosphino)ferrocene dichloropalladium (24 mg, 0.03 mmol), and sodium carbonate (2.5 mL of a 2.0 M aqueous solution, 5.00 mmol) were reacted to yield 2-(2-hydroxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 40 mg (15%) as a white solid: m.p. 305 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 2.86 (m, 2H), 3.46 (m, 2H), 6.92 (t, *J* = 7.5 Hz, 1H), 7.00 (d, *J* = 7.8 Hz, 1H), 7.16 (t, *J* = 7.8 Hz, 1H), 7.24 (m, 1H), 7.34 (dd, *J* = 7.5, 1.2 Hz, 1H), 7.55 (d, *J* = 7.8 Hz, 1H), 7.66 (d, *J* = 7.5 Hz, 1H), 8.00 (br t, 1H), 9.84 (br s, 1H), 11.20 (br s, 1H). MS (FAB, MH⁺): 279. Anal. (C₁₇H₁₄N₂O₂·0.44 CHCl₃) C, H, N.

15 Example EEE:

6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carbonitrile




Following a procedure from *JOC* 1998, p. 8224, 2-iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (100 mg, 0.32 mmol), sodium cyanide (31 mg, 0.64 mmol), palladium tetrakis(triphenylphosphine) (19 mg, 0.05 mmol), and copper(I) iodide were combined in a schlenk tube. The vessel was evacuated and refilled with

argon gas three times. Degassed propionitrile (2 mL) was added, and the reaction was stirred at 80 °C under an argon atmosphere for 15 h. The reaction mixture was partitioned between water and 25% iPrOH/CHCl₃. The layers were separated and the aqueous layer extracted thrice with 25% iPrOH/CHCl₃. The combined organic layers
5 were dried (MgSO₄) and concentrated in vacuo. The yellow solid was recrystallized from CH₂Cl₂/MeOH/hexanes to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carbonitrile, 38 mg (56%) as a pale yellow solid: m.p. 315 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.04 (m, 2H), 3.47 (m, 2H), 7.46 (t, *J* = 7.5 Hz, 1H), 7.64 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.81 (dd, *J* = 7.2, 0.9 Hz, 1H), 8.24 (br t, 1H), 12.44
10 (br s, 1H). MS (electrospray, [M+Na]⁺): 234. Anal. (C₁₂H₉N₃O) C, H, N.


Example FFF:

6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid octyl ester




15

Following a procedure similar to that described for Example MM (Compound 42), 2-iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (330 mg, 1.06 mmol), triethylamine (342 mg, 3.38 mmol), and palladium tetrakis(triphenylphosphine) (61 mg, 0.05 mmol) were reacted in 20 mL 1:1 n-octanol:DMF in a sealed tube under a
20 carbon monoxide atmosphere to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid octyl ester, 250 mg (58%), as a white solid: m.p. 170-171 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 0.85 (t, *J* = 7.2 Hz, 3H), 1.27 (m, 8H), 1.42 (m, 2H), 1.73 (m, 2H), 3.25 (m, 2H), 3.42 (m, 2H), 4.30 (t, *J* = 6.6 Hz, 3H), 7.38 (app t, *J* = 7.5 Hz, 1H), 7.62 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.74 (dd, *J* = 7.5, 0.9 Hz, 1H), 8.17 (br
25 t, 1H), 11.86 (br s, 1H). MS (FAB, MH⁺) 343. Anal. (C₂₀H₂₆N₂O₃) C, H, N.

Example GGG:**2-(4-Chloro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**

5 In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 4-chlorophenylboronic acid (195 mg, 1.24 mmol) were coupled to yield 2-(4-chloro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 223 mg (66%) as an off-white solid: m.p. 250-252 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.04 (m, 2H), 3.39 (m, 2H), 7.23 (app t, *J* = 7.5 Hz, 1H), 7.58 (m, 3H), 7.68 (m, 3H), 8.10 (br t, 1H), 11.66 (br s, 1H). MS (FAB, MH⁺) 297. Anal. (C₁₇H₁₃ClN₂O·0.8 H₂O) C, H, N.

10

Example HHH:**2-Pyridin-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**


15 In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 3-pyridylboronic acid (153 mg, 1.24 mmol) were coupled to yield 2-pyridin-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 75 mg (25%) as a light brown solid: m.p. 260.5-262.0 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.07 (m, 2H), 3.40 (m, 2H), 7.25 (app t, *J* = 7.8 Hz, 1H), 7.57 (m, 2H), 7.71 (dd, *J* = 7.5,

20

0.9 Hz, 1H), 8.05 (m, 1H), 8.12 (br t, 1H), 8.59 (m, 1H), 8.88 (m, 1H), 11.75 (br s, 1H). MS (FAB, MH⁺) 264. Anal. (C₁₆H₁₃N₃O·0.2 H₂O) C, H, N.

Example III:

5 **2-(2-Methoxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**




In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 2-methoxyphenylboronic acid (189 mg, 1.24 mmol) were coupled to yield 2-(2-methoxy-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 177 mg (53%) as a brown solid: m.p. 254-255 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.81 (m, 2H), 3.36 (m, 2H), 3.83 (s, 3H), 7.08 (app t, *J* = 7.5 Hz, 1H), 7.17 (m, 2H), 7.43 (m, 2H), 7.54 (dd, *J* = 7.8, 0.6 Hz, 1H), 7.67 (dd, *J* = 7.5, 0.6 Hz, 1H), 8.03 (br t, 1H), 11.27 (br s, 1H). MS (FAB, MH⁺) 293. Anal. (C₁₈H₁₆N₂O₂·0.3 H₂O) C, H, N.

15

Example JJJ:

2-Pyridin-4-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 4-pyridylboronic acid (153 mg, 1.24 mmol) were coupled to yield 2-pyridin-4-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 45 mg (15%)


20

as a beige solid: m.p. 250 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.13 (m, 2H), 3.41 (m, 2H), 7.29 (app t, *J* = 7.8 Hz, 1H), 7.63 (m, 3H), 7.72 (dd, *J* = 7.2, 0.9 Hz, 1H), 8.14 (br t, 1H), 8.69 (d, *J* = 6.0 Hz, 2H), 11.82 (br s, 1H). MS (FAB, MH⁺) 364. Anal. (C₁₆H₁₃N₃O) C, H, N.

5

Example KKK:


6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid sodium salt



10 In an attempt to form the piperazine amide, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methyl ester (100 mg, 0.41 mmol) was dissolved in 1 mL piperazine. The yellow solution was stirred under argon at 110 °C for 18 h. The reaction mixture was partitioned between saturated NaHCO₃ and 25% iPrOH/CHCl₃. The layers were separated and the aqueous layer extracted once with
15 25% iPrOH/CHCl₃. The combined organic layers were dried (MgSO₄) and concentrated in vacuo leaving ca. 3 mg of yellow solid. After standing overnight at room temperature, a pale yellow solid crystallized from the aqueous layer 80 mg (78%). The compound was identified as the sodium salt of 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid: m.p. 310 °C (dec); ¹H NMR (300
20 MHz, d₆-DMSO) δ 3.20 (m, 2H), 3.41 (m, 2H), 7.11 (app t, *J* = 7.8 Hz, 1H), 7.50 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.60 (dd, *J* = 7.5, 0.9 Hz, 1H), 7.96 (br t, 1H), 11.00 (br s, 1H). MS (electrospray, [M-Na]⁻) 229. Anal. (C₁₂H₉N₂O₃Na·0.5 H₂O) C, H, N.

Example LLL:


25 **2-(2-Methylsulfonyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**



In a manner similar to that described for Compound 12, the tricyclic bromide (530 mg, 2.00 mmol) and 2-thioanisole boronic acid (370 mg, 2.20 mmol) were coupled to yield 2-(2-methylsulfanyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 264 mg (43%) as an off-white solid: m.p. 271-272 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.39 (s, 3H), 2.73 (m, 2H), 3.37 (m, 2H), 7.23 (m, 2H), 7.37 (m, 2H), 7.49 (m, 2H), 7.70 (d, *J* = 7.2 Hz, 1H), 8.05 (br t, 1H), 11.41 (br s, 1H). MS (FAB, MH⁺) 309. Anal. (C₁₈H₁₆N₂OS) C, H, N.

10 Example MMM:


2-[4-(2-Pyrrolidin-1-yl-ethyl)-phenyl]-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for 2-[4-(2-diethylamino-ethyl)-phenyl]-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one (Example YY), the tricyclic bromide (198 mg, 0.75 mmol) and 1-[2-(4-bromo-phenyl)-ethyl]-pyrrolidine were coupled to yield 2-[4-(2-pyrrolidin-1-yl-ethyl)-phenyl]-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 160 mg (59%) as a beige solid: m.p. 228-229 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 1.69 (m, 4H), 2.51 (m, 4H), 2.67 (m, 2H), 2.81 (m, 2H), 3.05 (m, 2H), 3.39 (m, 2H), 7.20 (t, *J* = 7.8 Hz, 1H), 7.39 (d, *J* = 8.1 Hz, 2H), 7.56 (m, 3H), 7.68 (d, *J* = 7.5 Hz, 1H), 8.08 (br t, 1H), 11.31 (br s, 1H). MS (FAB, MH⁺): 360. Anal. (C₂₃H₂₅N₃O) C, H, N.

Example NNN:

N-[4-Fluoro-2-(6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-phenyl]-acetamide




5

In a manner similar to that described for 2-[4-(2-diethylamino-ethyl)-phenyl]-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one (Example YY), the tricyclic bromide (300 mg, 1.13 mmol) and *N*-(2-bromo-4-fluoro-phenyl)-acetamide (276 mg, 1.19 mmol) were coupled to yield *N*-[4-fluoro-2-(6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-phenyl]-acetamide, 83 mg (22%) as a beige solid: m.p. 260-261 °C (dec); ¹H NMR (300 MHz, *d*₆-DMSO) δ 1.97 (s, 3H), 2.66 (m, 2H), 3.33 (m, 2H), 7.25 (m, 3H), 7.56 (dd, *J* = 7.5, 0.6 Hz, 1H), 7.70 (dd, *J* = 7.2, 0.6 Hz, 1H), 7.76 (m, 1H), 8.04 (br t, 1H), 11.50 (br s, 1H). MS (FAB, MH⁺): 338. Anal. (C₁₆H₁₉FN₃O₂·0.16 H₂O) C, H, N.

15

Example OOO:

6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methylamide



20


6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methyl ester (50 mg, 0.20 mmol) was suspended in 1 mL of a 33% solution of

methylamine in methanol. The suspension was stirred at room temperature for 21 h. Another 2 mL 33% methylamine in methanol was added and the resulting solution stirred another 8 h at room temperature then 15 h at 30 °C. The reaction mixture was concentrated in vacuo leaving a yellow solid which was crystallized from

- 5 DMF/MeOH/CHCl₃ to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid methylamide, 36 mg (72%) as a yellow solid: m.p. 321-322 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 2.81 (s, 3H), 3.15 (m, 2H), 3.40 (m, 2H), 7.32 (app t, *J* = 7.8 Hz, 1H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.74 (d, *J* = 7.5 Hz, 1H), 7.95 (br q, 1H), 8.09 (br t, 1H), 11.46 (br s, 1H). MS (electrospray, [M+Na]⁺) 266. Anal.
- 10 (C₁₃H₁₃N₃O₂·0.4 H₂O) C, H, N.

Example PPP:

2-(4-Dimethylaminomethyl-3-fluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one




15

- 2-Fluoro-4-(6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-benzaldehyde (72 mg, 0.23 mmol. Prepared via the standard two-step, one-pot suzuki coupling of the tricyclic bromide and 4-bromo-2-fluoro-benzaldehyde as described for Example YY) was dissolved in 2 mL 2.0 M dimethylamine in
- 20 methanol. The orange solution was stirred at room temperature 10 min. The reaction was then cooled to 0 °C and a solution containing zinc chloride (17 mg, 0.13 mmol) and sodium cyanoborohydride (16 mg, 0.26 mmol) in 1 mL methanol, was added dropwise. The pH was adjusted to ca. 3 with concentrated HCl. The reaction was stirred for one hour as the temperature gradually warmed to room temperature. The
- 25 reaction was partitioned between CHCl₃ and water. The pH of the aqueous layer was adjusted to ca. 13 with solid KOH. The layers were separated, and the aqueous layer

extracted with 25% iPrOH/ CHCl₃. The combined organic layers were dried (MgSO₄) then concentrated in vacuo. Radial chromatography (eluting with 5% MeOH/ CHCl₃) then crystallization from CH₂Cl₂/hexanes yielded 2-(4-dimethylaminomethyl-3-fluoro-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 60 mg (76%) as a yellow solid.: m.p. 221.5-222.5 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.19 (s, 6H), 3.08 (m, 2H), 3.39 (m, 2H), 3.50 (s, 2H), 7.23 (app t, *J* = 7.8 Hz, 1H), 7.50 (m, 4H), 7.69 (d, *J* = 7.5 Hz, 1H) 8.10 (br t, 1H), 11.62 (br s, 1H). MS (FAB, MH⁺) 338. Anal. (C₂₀H₂₀FN₃O) C, H, N.

10 Example QQQ:


2-(3-Fluoro-4-pyrrolidin-1-ylmethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for as described for Example YY, the tricyclic bromide (1.00 g, 3.77 mmol) and 1-(4-bromo-2-fluoro-benzyl)-pyrrolidine (1.07 g, 4.19 mmol) were coupled to yield 2-(3-fluoro-4-pyrrolidin-1-ylmethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 150 mg (11%) as a beige solid: m.p. 139-140 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 1.71 (m, 4H), 2.50 (m, 4H, obscured by solvent), 3.07 (m, 2H), 3.40 (m, 2H), 3.68 (s, 2H), 7.23 (t, *J* = 7.8 Hz, 1H), 7.45 (m, 2H), 7.55 (m, 2H), 7.70 (dd, *J* = 7.5, 0.6 Hz, 1H), 8.07 (br t, 1H), 11.59 (br s, 1H). MS (electrospray, MH⁺) 364. Anal. (C₂₂H₂₂FN₃O·0.55 H₂O) C, H, N.

Example RRR:


25 2-Biphenyl-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and biphenyl-3-boronic acid (213 mg, 0.83 mmol) were coupled to yield 2-biphenyl-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 116 mg (30%) as an off-white crystalline solid: m.p. 160-163 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 3.13 (m, 2H), 3.42 (m, 2H), 7.24 (app t, *J* = 7.8 Hz, 1H), 7.42 (m, 1H), 7.61 (m, 7H), 7.79 (m, 2H), 7.94 (b s, 1H), 8.10 (br t, 1H), 11.67 (br s, 1H). MS (FAB, MH⁺) 339. Anal. (C₂₃H₁₈N₂O) C, H, N.

10 Example SSS:


2-(5-Chloro-2-methoxy-phenyl)-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Compound 12, the tricyclic bromide (129 mg, 0.49 mmol) and 5-chloro-2-methoxy-phenylboronic acid (100 mg, 0.54 mmol) were coupled to yield 2-(5-chloro-2-methoxy-phenyl)-3,4,5,6-tetrahydro-azepino[5,4,3-cd]indol-6-one, 100 mg (63%) as an off-white solid: m.p. 160-162 °C; ¹H NMR (300 MHz, d₆-DMSO) δ 2.81 (m, 2H), 3.34 (m, 2H), 3.84 (s, 3H), 7.20 (m, 2H), 7.46 (m, 2H), 7.55 (d, *J* = 7.8 Hz, 1H), 7.68 (d, *J* = 7.5 Hz, 1H), 8.05 (br t, 1H), 11.37 (br s, 1H). MS (FAB, MH⁺): 327. Anal. (C₁₈H₁₅ClN₂O₂) C, H, N, Cl.

Example TTT:


1,3,4,5,1',3',4',5'-Octahydro-[2,2']bi[azepino[5,4,3-cd]indolyl]-6,6'-dione




The title compound was isolated as a by-product of the coupling of the tricyclic bromide (642 mg, 2.42 mmol) under the conditions described for Example YY, 27 mg (6%) isolated as a yellow solid: m.p. <400 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 2.97 (m, 4H), 3.39 (m, 4H), 7.26 (t, *J* = 7.8 Hz, 2H), 7.59 (dd, *J* = 8.1, 0.9 Hz, 2H), 7.72 (dd, *J* = 7.5, 0.9 Hz, 2H), 8.12 (br t, 2H), 11.50 (br s, 2H). MS (electrospray, MH⁺): 372. Anal. (C₂₂H₁₈N₄O₂·0.25 H₂O) C, H, N.

10 Example UUU:


2-(3-Amino-phenylethynyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Example N, Compound 17, 3-ethynyl-aniline (129 mg, 1.10 mmol) was coupled to 2-iodo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (312 mmol, 1.00 mmol) to yield 2-(3-amino-phenylethynyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 250 mg (83%) as a pale yellow solid: m.p. 261-262 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.00 (m, 2H), 3.45 (m, 2H), 5.31 (br s, 2H), 6.63 (m, 1H), 6.71 (m, 1H), 6.76 (m, 1H), 7.08 (app t, *J* = 7.8 Hz, 1H), 7.26 (app t, *J* = 7.8 Hz, 1H), 7.48 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.70 (dd, *J* = 7.5, 0.6 Hz, 1H), 8.09 (br t, 1H), 11.75 (br s, 1H). MS (electrospray, MH⁺) 302. Anal. (C₁₉H₁₅N₃O·0.15 H₂O) C, H, N.

Example VVV:**2-(1H-Indol-5-yl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**

5 In a manner similar to that described for Compound 12, the tricyclic bromide (530 mg, 2.00 mmol) and indole-5-boronic acid (354 mg, 2.20 mmol) were coupled to yield 2-(1H-indol-5-yl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 396 mg (66%) as a beige solid: m.p. 315-317 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.10 (m, 2H), 3.41 (m, 2H), 6.54 (m, 1H), 7.17 (t, *J* = 7.8 Hz, 1H), 7.42 (m, 2H), 7.55 (m, 2H), 7.68 (d, *J* = 7.5 Hz, 1H), 7.83 (br s, 1H), 8.05 (br t, 1H), 11.26 (br s, 1H), 11.48 (br s, 1H). MS (electrospray, MH⁺) 302. Anal. (C₁₉H₁₃N₃O·0.25 H₂O) C, H, N.


Example WWW:**4-(6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-benzoic acid**

15 In a manner similar to that described for Compound 12, the tricyclic bromide (530 mg, 2.00 mmol) and 4-carboxyphenylboronic acid (365 mg, 2.20 mmol) were coupled to yield 4-(6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-benzoic acid, 340 mg (56%) as a pale yellow solid: m.p. 345.5-346.5 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.10 (m, 2H), 3.40 (m, 2H), 7.25 (t, *J* = 7.8 Hz, 1H), 7.59 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.70 (dd, *J* = 7.5, 0.6 Hz, 1H), 7.78 (m, 2H), 8.10 (m,

3H), 11.73 (br s, 1H), 13.00 (br s, 1H). MS (electrospray, MH⁺) 307. Anal. (C₁₈H₁₄N₂O₃·0.9 H₂O) C, H, N.

Example XXX:

5 **6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid**




6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid octyl ester (Example FFF) (350 mg, 1.02 mmol) and lithium hydroxide (122 mg, 5.11 mmol) were dissolved in 10 mL 2:1 methanol:water and stirred at room temperature for 24 h.

- 10 The reaction mixture was diluted with water then washed twice with dichloromethane. The aqueous solution was acidified to ca. pH 2 with conc. HCl. The white precipitate was collected by filtration, washed with water, and dried in vacuo to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid, 235 mg (99%) as a white solid: m.p. 298-299 °C (dec); ¹H NMR (300 MHz, d₆-
- 15 DMSO) δ 3.17 (m, 2H), 3.41 (m, 2H), 7.35 (t, *J* = 7.8 Hz, 1H), 7.59 (d, *J* = 8.1 Hz, 1H), 7.73 (d, *J* = 7.5 Hz, 1H), 8.14 (br t, 1H), 11.77 (br s, 1H), 13.14 (br s, 1H). MS (electrospray, MH⁺): 231. Anal. (C₁₂H₁₀N₂O₃·1.0 H₂O) C, H, N.

Example YYY:


- 20 **6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (4-fluorophenyl)-amide**



- 6-Oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (100 mg, 0.43 mmol), 4-fluoroaniline (48 mg, 0.43 mmol), and diisopropylethylamine (168 mg, 1.30 mmol) were dissolved in 5 mL dry DMF. HATU (173 mg, 0.46 mmol) was added and the resulting mixture stirred at room temperature under argon for 3 d. The reaction mixture was partitioned between water and 25% iPrOH/CHCl₃. The layers were separated, and the aqueous layer extracted thrice with 25% iPrOH/CHCl₃. The combined organic layers were dried (MgSO₄) and concentrated in vacuo leaving an off-white solid which was recrystallized from chloroform/methanol to yield 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (4-fluorophenyl)-amide, 70 mg (50%) as a pale yellow solid: m.p. 330-332 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 3.28 (m, 2H), 3.42 (m, 2H), 7.22 (m, 2H), 7.35 (app t, *J* = 7.8 Hz, 1H), 7.65 (dd, *J* = 7.8, 0.6 Hz, 1H), 7.77 (m, 3H), 8.16 (br t, 1H), 10.08 (br s, 1H), 11.81 (br s, 1H). MS (electrospray, MH⁺) 324. Anal. (C₁₈H₁₄FN₃O₂·0.4 H₂O) C, H, N.

Example ZZZ:

(4-Chloro-phenyl)-1,5-dihydro-[1,2]diazepino[4,5,6-cd]indol-6-one



- 2-Iodo-3-nitro-benzoic acid methyl ester:** 2-iodo-3-nitro-benzoic acid (61 g, 208 mmol, prepared as described in *Org. Syn. Coll. Vol. I*, 56-58, and 125-127), sulfuric acid (40.8 g, 416 mmol), and trimethyl orthoformate (88.4 g, 833 mmol) were dissolved in 500 mL dry MeOH. The reaction was refluxed under argon for 20 h. The reaction mixture was concentrated to 100 mL then partitioned between saturated $\text{NaHCO}_{3(\text{aq})}$ and CH_2Cl_2 . The layers were separated and the aqueous layer
- 10 extrated three times with CH_2Cl_2 . The combined organic layers were dried (MgSO_4) and concentrated in vacuo. The yellow solid was crystallized from CH_2Cl_2 /hexanes yielding 2-iodo-3-nitro-benzoic acid methyl ester, 57.8 g (90%) as a yellow solid: m.p. 64.0-64.5 °C; ^1H NMR (300 MHz, CDCl_3) 3.99 (s, 3H), 7.54 (app t, $J = 7.8$ Hz, 1H), 7.70 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.77 (dd, $J = 7.8, 1.8$ Hz, 1H).
- 15 **3-Amino-2-iodo-benzoic acid methyl ester:** 2-Iodo-3-nitro-benzoic acid methyl ester (1.00 g, 3.26 mmol) was dissolved in 15 mL MeOH. Tin (II) chloride (2.78 g, 14.66 mmol) and water (0.35 g, 19.54 mmol) were added and the yellow

solution stirred at room temperature for 20 h. Celite was added to the solution followed by 10 mL 3 M NaOH. The suspension was diluted with MeOH and the precipitate filtered off. The filter cake was washed with three portions boiling CH_2Cl_2 . The layers were separated and the aqueous layers extracted once with CH_2Cl_2 . The combined organic layers were dried (MgSO_4) and concentrated in vacuo to yield 2-iodo-3-nitro-benzoic acid methyl ester, 0.89 g (99%), as a clear oil. ^1H NMR (300 MHz, d_6 -DMSO) 3.81 (s, 3H), 5.52 (br s, 2H), 6.72 (dd, $J = 7.5, 1.2$ Hz, 1H), 6.87 (dd, $J = 7.5, 1.2$ Hz, 1H), 6.87 (dd, $J = 7.8, 1.2$ Hz, 1H), 7.12 (app t, $J = 7.5$ Hz, 1H). MS (electrospray, MH^+) 278.

10 **3-Amino-2-(4-chloro-phenylethynyl)-benzoic acid methyl ester:** 2-iodo-3-nitro-benzoic acid methyl ester (0.79 g, 2.84 mmol), 1-chloro-4-ethynylbenzene (0.41 g, 2.99 mmol), palladium tetrakis(triphenylphosphine) (0.16 g, 0.14 mmol), copper (I) iodide (0.03 g, 0.14 mmol), and triethylamine (1.44 g, 14.19 mmol) were dissolved in 15 mL toluene. Argon was bubbled through the resulting solution for 15 min. The reaction was stirred under argon at 80 °C for 2 h and 20 min. The reaction mixture was then washed once with water, dried (MgSO_4), and concentrated in vacuo. The orange oil was purified by flash chromatography eluting with 50 to 100% CHCl_3 /hexanes to yield 3-amino-2-(4-chloro-phenylethynyl)-benzoic acid methyl ester, 0.76 g (94%) as a yellow oil. ^1H NMR (300 MHz, d_6 -DMSO) 3.84 (s, 3H), 5.84 (br s, 2H), 6.97 (dd, $J = 8.1, 1.3$ Hz, 1H), 7.05 (dd, $J = 7.5, 1.2$ Hz, 1H), 7.17 (app t, $J = 7.5$ Hz, 1H), 7.49 (d, $J = 8.7$ Hz, 2H), 7.63 (d, $J = 8.7$ Hz, 2H). MS (electrospray, MH^+) 286.

25 **2-(4-Chloro-phenyl)-1H-indole-4-carboxylic acid methyl ester:** 3-amino-2-(4-chloro-phenylethynyl)-benzoic acid methyl ester (0.73 g, 2.54 mmol) and palladium (II) chloride (23 mg, 0.13 mmol) were combined in 10 mL acetonitrile. The yellow solution was stirred under argon at 75 °C for 17 h. The solvent was removed in vacuo leaving an orange solid which was purified by flash chromatography eluting with 50 to 100% CH_2Cl_2 /hexanes. 2-(4-chloro-phenyl)-1H-indole-4-carboxylic acid methyl ester; 0.53 g (72%) was isolated as an off-white solid: m.p. 150.0-151.5 °C; ^1H NMR (300 MHz, d_6 -DMSO) 3.93 (s, 3H), 7.23 (app

30

t, $J = 7.8$ Hz, 1H), 7.41 (d, $J = 1.8$ Hz, 1H), 7.57 (d, $J = 8.7$ Hz, 2H), 7.68 (d, $J = 7.8$ Hz, 1H), 7.75 (dd, $J = 7.5, 0.9$ Hz, 1H), 7.95 (d, $J = 8.7$ Hz, 2H), 11.99 (br s, 1H).

HRMS (MALDI, MH^+) Calcd for $C_{16}H_{12}ClNO_2$: 286.0635. Found: 286.0631.


2-(4-Chloro-phenyl)-3-formyl-1H-indole-4-carboxylic acid methyl ester:

5 Phosphorous oxychloride (0.42 g, 2.71 mmol) was added to DMF (0.99 g, 13.57 mmol) at 0 °C. The resulting colorless solution was added dropwise to a solution of 2-(4-chloro-phenyl)-1H-indole-4-carboxylic acid methyl ester (0.52 g, 1.81 mmol) in 10 mL dry CH_2Cl_2 at 0 °C. The reaction was stirred at 0 °C for 10 min then quenched by addition of 5 mL 2 M NaOAc_(aq). The layers were separated and the aqueous layer
10 extracted once with CH_2Cl_2 . The combined organic layers were dried ($MgSO_4$) then concentrated in vacuo leaving an orange oil which crystallized on standing. The crystals were rinsed with CH_2Cl_2 then dried in vacuo to yield 2-(4-chloro-phenyl)-3-formyl-1H-indole-4-carboxylic acid methyl ester, 231 mg (41%) as an off-white solid: m.p. 221-222 °C; 1H NMR (300 MHz, d_6 -DMSO) 3.93 (s, 3H), 7.49 (app t, J
15 = 7.5 Hz, 1H), 7.71 (m, 4H), 7.94 (d, $J = 7.8$ Hz, 2H), 9.71 (s, 1H), 13.67 (br s, 1H). MS (electrospray, $[M-H]$) 312.

(4-Chloro-phenyl)-1,5-dihydro-[1,2]diazepino[4,5,6-cd]indol-6-one: 2-(4-chloro-phenyl)-3-formyl-1H-indole-4-carboxylic acid methyl ester (100 mg, 0.32 mmol) was dissolved in 5 mL MeOH. Hydrazine (30 mg, 0.92 mmol) was added
20 causing the immediate precipitate. Acetic acid (13 mg, 0.22 mmol) was added and the yellow suspension refluxed for 1.5 h. The yellow solid was collected by filtration, rinsed once with MeOH, then dried in vacuo to give (4-chloro-phenyl)-1,5-dihydro-[1,2]diazepino[4,5,6-cd]indol-6-one, 55 mg (59%) as a bright yellow solid: m.p. 324.0-324.5 °C (dec); 1H NMR (300 MHz, d_6 -DMSO) 7.23 (app t, $J = 7.8$ Hz,
25 1H), 7.49 (s, 1H), 7.55 (m, 2H), 7.65 (d, $J = 8.7$ Hz, 2H), 7.71 (d, $J =$ Hz, 2H), 10.36 (s, 1H), 12.32 (br s, 1H). HRMS (MALDI, MH^+) Calcd for $C_{16}H_{10}ClN_3O$: 296.0591. Found: 296.0586. Anal. ($C_{16}H_{10}ClN_3O \cdot 0.5 H_2O$) C, H, N.

Example AAAA:


30 2-(4-Fluoro-phenyl)-1,5-dihydro-[1,2]diazepino[4,5,6-cd]indol-6-one



In a manner similar to that described for Example ZZZ, 2-(4-fluorophenyl)-3-formyl-1H-indole-4-carboxylic acid methyl ester (145 mg, 0.49 mmol) was
 5 condensed with hydrazine (45 mg, 1.41 mmol) to give 2-(4-fluorophenyl)-1,5-dihydro-[1,2]diazepino[4,5,6-cd]indol-6-one, 120 mg (88%) as a bright yellow solid: m.p. 340-341 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 7.22 (app t, *J* = 7.8 Hz 1H), 7.43 (m, 3H), 7.54 (m, 2H), 7.73 (m, 2H), 10.33 (s, 1H), 12.23 (br s, 1H). MS
 10 (electrospray, MH⁺) 280. Anal. (C₁₆H₁₀FN₃O) C, H, N.

Example BBBB:

2-Thiophen-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one




15

In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and thiophene-2-boronic acid (159 mg, 1.24 mmol) were
 coupled to yield 2-thiophen-2-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one,
 20 171 mg (56%) as a beige solid: m.p. 220.5-222.5 °C; ¹H NMR (300 MHz d₆-DMSO) 3.08 (m, 2H), 3.48 (m, 2H), 7.23 (m, 2H), 7.52 (m, 2H), 7.69 (m, 2H), 8.05 (br t,

1H), 11.60 (br s, 1H). MS (electrospray, MH⁺) 269. Anal. (C₁₅H₁₂N₂)S·0.8 H₂O) C, H, N.

Example CCCC:


5 **2-Thiophen-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**



In a manner similar to that described for Compound 12, the tricyclic bromide
 10 (300 mg, 1.13 mmol) and thiophene-3-boronic acid (159 mg, 1.24 mmol) were
 coupled to yield 2-thiophen-3-yl-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one,
 249 mg (82%) as a beige solid: m.p. 255-256 °C; ¹H NMR (300 MHz, d₆-DMSO)
 3.08 (m, 2H), 3.43 (m, 2H), 7.19 (t, *J* = 7.8 Hz, 1H), 7.54 (m, 2H), 7.67 (dd, *J* = 7.5,
 0.9 Hz, 1H), 7.74 (m, 1H), 7.78 (m, 1H), 8.03 (br t, 1H), 11.49 (br s, 1H). MS
 15 (electrospray, MH⁺) 269. Anal. (C₁₅H₁₂N₂OS·0.35 H₂O) C, H, N, S.

Example DDDD:

2-(1H-Pyrrol-2-yl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



20


In a manner similar to that described for Compound 12, the tricyclic bromide (300 mg, 1.13 mmol) and 1-(t-butoxycarbonyl)pyrrole-2-boronic acid (263 mg, 1.24 mmol) were coupled with concomitant removal of the BOC group to yield 2-(1H-pyrrol-2-yl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 81 mg (28% as a

5 greenish grey solid: m.p. >400 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 3.02 (m, 2H), 3.42 (m, 2H), 6.22 (m, 1H), 6.44 (m, 1H), 6.97 (m, 1H), 7.14 (t, *J* = 7.5 Hz, 1H), 7.49 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.64 (dd, *J* = 7.5, 0.6 Hz, 1H), 7.98 (br t, 1H), 11.01 (br s, 1H), 11.13 (br s, 1H). MS (electrospray, MH⁺) 252. Anal. (C₁₅H₁₃N₃O·0.4 H₂O) C, H, N.

10

Example EEEE:


2-(4-Methylsulfanyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



15

In a manner similar to that described for Compound 12, the tricyclic bromide (1.00 g, 3.77 mmol) and 4-thioanisole boronic acid (0.70 g, 4.15 mmol) were coupled to yield 2-(4-methylsulfanyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 416 mg (36%) as a beige solid: m.p. 250-251 °C; ¹H NMR (300 MHz, d₅-DMSO)


20 2.54 (s, 3H), 3.03 (m, 2H), 3.39 (m, 2H), 7.20 (t, *J* = 7.8 Hz, 1H), 7.41 (d, *J* = 7.5, 0.9 Hz, 1H), 8.04 (br t, 1H), 11.52 (br s, 1H). MS (electrospray, MH⁺) 309. Anal. (C₁₈H₁₆N₂OS·0.6 H₂O) C, H, N.

Example FFFF:**2-(4-Methanesulfinyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**

5

2-(4-methylsulfonyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one (100 mg, 0.32 mmol) was dissolved in 10 mL 1:1 MeOH:CH₂Cl₂. The solution was cooled to 0 °C and oxone (259 mg, 0.42 mmol) was added dropwise as a solution in 1.5 mL H₂O). The bright yellow reaction mixture was stirred at 0 °C for 15 min.

- 10 Saturated Na₂S₂O_{5(aq)} (4mL) was added. The layers were separated and the aqueous layer extracted twice with 25% iPrPH/CHCl₃. The combined organic layers were dried (MgSO₄), concentrated in vacuo, and the two products (2-(4-methanesulfinyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one and 2-(4-methanesulfonyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one) separated by radial
- 15 chromatography eluting with 5% MeOH/CHCl₃. Each was then crystallized from CH₂Cl₂/MeOH. 2-(4-Methanesulfinyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 39 mg (37%), was isolated as a white solid: m.p. 316-317 °C (dec);
- ¹H NMR (300 MHz, d₆-DMSO) 2.81 (s, 3H), 3.09 (m, 2H), 3.40 (m, 2H), 7.25 (t, *J* = 7.8 Hz, 1H), 7.59 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.71 (dd, *J* = 7.5, 0.9 Hz, 1H), 7.84 (m,
- 20 4H), 8.08 (br t, 1H), 11.68 (br s, 1H). MS (electrospray, MH⁺) 325. Anal. (C₁₈H₁₆N₂O₂S) C, H, N, S.

Example GGGG:**2-(4-Methanesulfonyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**


5

2-(4-methanesulfonyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 20 mg (18%) was isolated in the chromatography described above as a white solid: m.p. 308-309 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 3.10 (m, 2H), 3.28 (s, 3H), 3.41 (m, 2H), 7.28 (t, *J* = 7.8 Hz, 1H), 7.61 (dd, *J* = 8.1, 0.6 Hz, 1H), 7.72 (dd, *J* = 7.5, 0.6 Hz, 1H), 7.91 (d, *J* = 8.4 Hz, 2H), 8.06 (d, *J* = 8.4 Hz, 2H), 8.11 (br t, 1H), 11.77 (br s, 1H). MS (electrospray, MH⁺) 341. Anal. (C₁₈H₁₆N₂O₃S) C, H, N, S.

10

Example HHHH:**2-Bromo-8-fluoro-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**

15




The title compound was prepared in a manner similar to that used for 2-bromo-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, starting from 5-fluoro-2-methylbenzoic acid. 2-Bromo-8-fluoro-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one was isolated as an orange solid: m.p. 203-204 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) δ 2.79 (m, 2H), 3.41 (m, 2H), 7.29 (dd, *J* = 8.7, 1.2 Hz, 1H), 7.74 (dd, *J* =

20

10.8, 1.5 Hz, 1H), 8.23 (br t, 1H), 12.12 (br s, 1H). MS (electrospray, $[M+Na]^+$) 305/307.


5 **8-Fluoro-2-(3-methylaminomethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**



3-(8-fluoro-6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-
 10 benzaldehyde (247 mg, 0.80 mmol; prepared in a manner similar to that described for compound 12 from 2-bromo-8-fluoro-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one and 3-formylphenylboronic acid) was reacted with methylamine (4.91 mmol) as described for Compound PPP to yield 8-fluoro-2-(3-methylaminomethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 193 mg (74%) as an off-white solid:
 15 m.p. 270-272 °C (dec); ^1H NMR (300 MHz, d_6 -DMSO) 2.34 (s, 3H), 3.05 (m, 2H), 3.39 (m, 2H), 3.78 (s, 2H), 7.42 (m, 5H), 7.61 (br s, 1H), 8.26 (br t, 1H), 11.70 (br s, 1H). HRMS (MALDI, MH^+) Calcd for $C_{19}H_{18}N_3OF$: 324.1512. Found: 324.1498. Anal. ($C_{19}H_{18}N_3OF \cdot 1.5 H_2O \cdot 0.35 CHCl_3$) C, H, N.

20 **Example III:**


8-Fluoro-2-(4-methylaminomethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one



4-(8-fluoro-6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl)-
benzaldehyde (100 mg, 0.32 mmol; prepared in a manner similar to that described for
5 compound 12, for 2-bromo-8-fluoro-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one
and 4-formylphenylboronic acid) was reacted with methylamine (1.62 mmol) as
described for Compound PPP to yield 8-fluoro-2-(4-methylaminomethyl-phenyl)-
1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 32 mg (31%) as a yellow solid:
m.p. 1543-155 °C; ¹H NMR (300 MHz, d₆-DMSO) 2.28 (s, 3H), 3.04 (m, 2H), 3.40
10 (m, 2H), 3.69 (s, 2H), 7.32 (dd, *J* = 9.0, 2.4 Hz, 1H), 7.44 (m, 3H), 7.57 (d, *J* = 8.1
Hz, 2H), 8.25 (br t, 1H), 11.67 (br s, 1H). HRMS (MALDI MH⁺) Calcd for
C₁₉H₁₈N₃OF: 324,1512. Found: 325.1524. Anal. (C₁₉H₁₈N₃OF·0.3 H₂O) C, H, N.

Example JJJJ:

15 **8-Fluoro-2-(4-pyrrolidin-1-ylmethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one**




20 In a manner similar to that described for Compound PPP, 4-(8-fluoro-6-oxo-
3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-2-yl) benzaldehyde (100 mg, 0.32
mmol; prepared in a manner similar to that described for compound 12 from 2-

bromo-8-fluoro-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one and 4-formylphenylboronic acid) was reacted with pyrrolidine (115 mg, 1.62 mmol) to yield 8-fluoro-2-(4-pyrrolidin-1-ylmethyl-phenyl)-1,3,4,5-tetrahydro-azepino[5,4,3-cd]indol-6-one, 16 mg (14%) as a yellow solid: m.p. 264-265 °C (dec), ¹H NMR (300 MHz, d₆-DMSO) 1.72 (m, 4H), 2.49 (m, 4H), 3.04 (m, 2H), 3.39 (m, 2H), 3.64 (br s, 2H), 7.31 (dd, *J* = 9.3, 2.4 Hz, 1H), 7.43 (m, 3H), 7.58 (d, *J* = 8.1 Hz, 2H), 8.25 (br t, 1H), 11.66 (br s, 1H). HRMS (MALDI MH⁺) Calcd for C₂₂H₂₂N₃OF: 362.1825. Found: 364.1810. Anal. (C₂₂H₂₂N₃OF·0.5 H₂O) C, H, N.

10 Example KKKK:

6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid phenylamide




15

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with aniline (27 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid phenylamide as a white solid: m.p. 320-322 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 3.28 (m, 2H), 3.42 (m, 2H), 7.11 (app t, *J* = 7.5 Hz, 1H), 7.37 (m, 3H), 7.64 (d, *J* = 8.1 Hz, 1H), 7.74 (m, 3H), 8.15 (br t, 1H), 9.98 (br s, 1H), 11.78 (br s, 1H). MS (electrospray, MH⁺) 306. Anal. (C₁₈H₁₅N₃O₂·0.25 H₂O) C, H, N.

Example LLLL:

6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (4-chloro-phenyl)-amide




5

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with 4-chloroaniline (37 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (4-chloro-phenyl)-amide as a white solid: ^1H NMR (300 MHz, d_6 -DMSO) 3.26 (m, 2H), 3.42 (m, 2H), 7.36 (app t, J = 7.8 Hz, 1H), 7.44 (d, J = 8.7 Hz, 2H), 7.65 (d, J = 8.1 Hz, 1H), 7.76 (m, 3H), 8.16 (br t, 1H), 10.12 (br s, 1H), 11.79 (br s, 1H). MS (electrospray, MH^+) 340. Anal. ($\text{C}_{18}\text{H}_{14}\text{ClN}_3\text{O}_2$) C, H, N.

15

Example MMMM:

6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid naphthalen-2-ylamide




20

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-zepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with 2-naphthylamine (41 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid naphthalen-2-ylamide as a white solid:

- 5 ¹H NMR (300 MHz, d₆-DMSO) 3.33 (m, 2H), 3.45 (m, 2H), 7.38 (app t, *J* = 7.8 Hz, 1H), 7.47 (m, 2H), 7.68 (d, *J* = 8.1 Hz, 1H), 7.78 (m, 2H), 7.91 (m, 3H), 8.19 (br t, 1H), 8.43 (br s, 1H), 10.21 (br s, 1H), 11.84 (br s, 1H). MS (electrospray, MH⁺) 356. Anal. (C₂₂H₁₇N₃O₂·0.7 H₂O) C, H, N.

10 Example NNNN:


6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid naphthalen-1-ylamide



15


In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with 1-naphthylamine (41 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid naphthalen-1-ylamide as a white solid:

- 20 m.p. 330-332 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 3.33 (m, 2H), 3.48 (m, 2H), 7.38 (app t, *J* = 7.8 Hz, 1H), 7.57 (m, 3H), 7.68 (d, *J* = 7.8 Hz, 1H), 7.77 (m, 2H), 7.87 (d, *J* = 7.8 Hz, 1H), 7.99 (m, 1H), 8.13 (m, 2H), 10.06 (br s, 1H), 11.87 (br s, 1H). MS (electrospray, MH⁺) 356. Anal. (C₂₂H₁₇N₃O₂·0.5 H₂O) C, H, N.

Example OOOO:**6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid prop-2-ynylamide**

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with propargylamine (16 mg; 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid prop-2-ynylamide as a white solid: m.p. 191-192 °C; ¹H NMR (300 MHz, d₆-DMSO) 3.19 (m, 3H), 3.39 (m, 2H), 4.10 (m, 2H), 7.32 (app t, *J* = 7.8 Hz, 1H), 7.59 (d, *J* = 8.1 Hz, 1H), 7.72 (d, *J* = 7.2 Hz, 1H), 8.12 (br t, 1H), 8.43 (br t, 1H), 11.60 (br s, 1H). MS (electrospray, MH⁺) 268. Anal. (C₁₅H₁₃N₃O₂·2H₂O) C, H, N.

15


Example PPPP:**6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid isopropyl-amide**

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with isopropylamine (17 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid isopropylamide as a white solid: m.p.

5 261-262 °C (dec); ¹H NMR (300 MHz, d₆-DMSO) 1.20 (d, *J* = 6.6 Hz, 1H) 3.22 (m, 2H), 3.38 (m, 2H), 4.90 (m, 1H), 7.32 (app t, *J* = 7.8 Hz, 1H), 7.59 (d, *J* = 8.1 Hz, 1H), 7.71 (d, *J* = 7.2 Hz, 1H), 7.81 (d, *J* = 7.5 Hz, 1H), 8.10 (br t, 1H), 11.53 (br s, 1H). MS (electrospray, MH⁺) 272. Anal. (C₁₅H₁₇N₃O₂·0.2H₂O) C, H, N.

10 Example QQQQ:

6-Oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid cyclopropylamide




15

In a manner similar to that described for Compound YYY, 6-oxo-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid (60 mg, 0.26 mmol) was coupled with cyclopropylamine (17 mg, 0.29 mmol) to yield 6-oxo-1,3,4,5-tetrahydro-1H-azepino[5,4,3-cd]indole-2-carboxylic acid cyclopropyleamide as a

20 white solid: m.p. 249-251 °C; ¹H NMR (300 MHz, d₆-DMSO) 0.56 (m, 2H), 0.75 (m, 2H), 2.95 (m, 2H), 3.37 (m, 2H), 3.61 (m, 1H), 7.30 (app t, *J* = 7.5 Hz, 1H), 7.58 (d, *J* = 8.1 Hz, 1H), 7.70 (d, *J* = 7.2 Hz, 1H), 8.09 (m, 2H), 11.48 (br s, 1H). MS (electrospray, MH⁺) 270. Anal. (C₁₅H₁₅N₃O₂·1H₂O) C, H, N.

Example RRRR:

(rac)-3-(4-Methoxyphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one




5

In a manner similar to that described for the preparation of Example Q, methyl indole-4-carboxylate and p-methoxy nitrostyrene were condensed and the resulting nitro alkane was reduced/cyclized to give, after recrystallization

(CH₂Cl₂/MeOH/hexanes), (rac)-3-(4-methoxyphenyl)-3,4,5,6-tetrahydro-1H-
10 axepino[5,4,3-cd]indol-6-one, 16.9 mg (50%) as a white solid: m.p. 221-223 °C; ¹H
NMR (300 MHz, d₄-MeOH) 3.57 (br m, 5H), 5.15 (br s, 1H) 6.62 (m, 2H), 6.86 (m,
2H), 7.08 (app t, J = 7.8 Hz, 1H), 7.11 (s, 1H), 7.37 (d, J = 7.9 Hz, 1H), 7.73 (d, J =
7.5 Hz, 1H). Anal. (C₁₉H₁₆N₂O₂·0.25 H₂O) C, H, N.

15 Example SSSS:

2-(3-Morpholin-4-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-
6-one




20

In a manner similar to that described for Compound 22, the aldehyde 15 (29 mg, 0.1 mmol) in MeOH (1 mL) was treated with morpholine (0.04 mL, 0.5 mmol) and a solution of sodium cyanoborohydride (0.15 mmol) and zinc chloride (0.08 mmol) in MeOH (1 mL) to give, after radial chromatography (5% MeOH in CHCl₃),
 5 2-(3-morpholin-4-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 35 mg (99%) as sticky white solid: ¹H NMR (300 MHz, d₆-DMSO) 2.37 (m, 4H), 3.02 (m, 2H), 3.35 (m, 2H), 3.51 (m, 6H), 7.17 (app t, *J* = 7.7 Hz, 1H), 7.30 (br d, 1H), 7.52 (m, 4H), 7.64 (d, *J* = 7.5 Hz, 1H), 8.03 (br t, 1H), 11.53 (br s, 1H). HRMS (FAB, MH⁺) Calcd for C₂₂H₂₄N₃O₂: 362.1869. Found: 362.1866.

10

Compound TTTT:

2-(3-Pyrrolidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one




15

In a manner similar to that described for Compound 22, the aldehyde 15 (200 mg, 0.69 mmol) in MeOH (10 mL) was treated with pyrrolidine (0.34 mL, 4.14 mmol) and a solution of sodium cyanoborohydride (0.76 mmol) and zinc chloride
 20 (0.38 mmol) in MeOH (1.4 mL) to give, after crystallization (CH₂Cl₂/MeOH/hexanes), 2-(3-pyrrolidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 139 mg (58%) as pale yellow solid: m.p. 219-223 °C (dec); ¹H NMR (300 MHz, d⁶-DMSO) 1.73 (m, 4H), 2.49 (m, 4H), 3.06 (m, 2H), 3.40 (m, 2H), 3.69 (s, 2H), 7.22 (t, *J* = 7.7 Hz, 1H), 7.34 (br d, 1H), 7.53 (m, 4H),
 25 7.68 (dd, *J* = 7.7, 0.8 Hz, 1H), 8.08 (br t, 1H), 11.59 (br s, 1H). HRMS (FAB, MH⁺)

Calcd for $C_{22}H_{24}N_3O$: 346.1919. Found: 346.1910. Anal. ($C_{22}H_{25}N_3O \cdot 0.6 H_2O$) C, H, N.

Example UUUU:


2-(4-Pyrrolidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one



In a manner similar that described for Compound 22, the para-aldehyde (150 mg, 0.52 mmol) in MeOH (10 mL) was treated with pyrrolidine (0.26 mL, 3.10 mmol) and a solution of sodium cyanoborohydride (0.57 mmol) and zinc chloride (0.28 mmol) in MeOH (1.1 mL) to give, after crystallization (CH_2Cl_2 /MeOH/hexanes), 2-(4-pyrrolidin-1-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 141 mg (79%) as pale yellow solid: m.p. 221-225 °C (dec); 1H NMR (300 MHz, d_6 -DMSO) 1.71 (m, 4H), 2.46 (m, 4H), 3.06 (m, 2H), 3.41 (m, 2H), 3.41 (m, 2H), 3.63 (s, 2H), 7.21 (t, $J = 7.8$ Hz, 2H), 7.45 (d of Abq, $J = 8.2$ Hz, 2H), 7.55 (dd, $J = 7.9, 0.9$ Hz, 1H), 7.59 (d of Abq, $J = 8.2$ Hz, 2H), 7.68 (br d, 1H), 8.07 (br t, 1H), 11.54 (br s, 1H). HRMS (FAB, MH^+) Calcd for $C_{22}H_{24}N_3O$: 346.1919. Found: 346.1911. Anal. ($C_{22}H_{25}N_3O \cdot 0.5 H_2O$) C, H, N.

Example VVVV:


2-(4-Morpholin-4-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one



In a manner similar to that described for Compound 22, the para-aldehyde (264 mg, 0.91 mmol) in MeOH (10 mL) was treated with morpholine (0.40 mL, 4.55 mmol) and a solution of sodium cyanoborohydride (1.36 mmol) and zinc chloride (0.68 mmol) in MeOH (2.0 mL) to give, after recrystallization (CH₂Cl₂/MeOH/hexanes) and radial chromatography, 2-(4-morpholin-4-ylmethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 44.8 mg (14%) as solid: ¹H NMR (300 MHz, d₆-DMSO) 2.39 (m, 4H), 3.06 (m, 2H), 3.41 (m, 2H), 3.53 (s, 2H), 3.59 (m, 4H), 7.21 (br t, 1H), 7.46 (d of Abq, *J* = 8.0 Hz, 2H), 7.55 (br d, 1H), 7.62 (d of Abq, *J* = 8.0 Hz, 2H), 7.68 (br d, 1H), 8.07 (br t, 1H), 11.55 (br s, 1H). HRMS (FAB, MH⁺) Calcd for C₂₂H₂₄N₃O₂: 362.1869. Found: 362.1861.

Example WWW:

15 2-(4-Hydroxymethylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one




20 The title compound was isolated as a reduction by-product from the reductive amination of the para-paldehyde with morpholine and sodium cyanoborohydride, and recrystallized (CH₂Cl₂/MeOH/hexanes) to give 2-(4-hydroxymethylphenyl)-3,4,5,6-


tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, 64 mg (24%) as a white solid: ^1H NMR (300 MHz, d_6 -DMSO) 3.05 (m, 2H), 3.39 (m, 2H), 4.57 (d, $J = 5.6$ Hz, 2H), 5.27 (t, $J = 5.6$ Hz, -OH), 7.21 (br t, 1H), 7.47 (d of Abq, $J = 7.9$ Hz, 2H), 7.55 (br d, 1H), 7.62 (d of Abq, $J = 7/9$ Hz, 2H), 7.68 (br d, 1H), 8.07 (br t, 1H), 11.55 (s, 1H). Anal. (C₁₈H₁₆N₂O₂·0.9 H₂O) C, H, N.

Example XXXX:

2-(4-(N,N-Dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, N-oxide



A solution of Compound 21 (58 mg) in acetone (7.0 mL) was treated with 30% aqueous hydrogen peroxide (0.6 mL) at room temperature and the yellow solution was allowed to stir for three days. The acetone was removed in vacuo and the residue was taken-up in isopropyl alcohol. A solid was precipitated with the addition of an equal volume of cold hexanes and collected by a quick filtration. Precautions were taken to prevent the solid from absorbing moisture from the atmosphere. The solid was recrystallized (isopropanol/acetone/CH₂Cl₂/hexanes) to give 2-(4-(N,N-dimethylamino)methylphenyl)-3,4,5,6-tetrahydro-1H-azepino[5,4,3-cd]indol-6-one, N-oxide, 37 mg (60%) as a pale yellow solid: ^1H NMR (300 MHz, d_6 -DMSO) 3.22 (s, 6H), 3.56 (br m, 4H), 4.63 (s, 2H), 7.40 (br t, 1H), 7.76 (br d, 1H), 7.87 (m, 5H), 8.29 (br t, 1H), 12.00 (br s, 1H). HRMS (FAB, $\text{MH}^+ - \text{H}_2\text{O}$) Calcd for C₂₀H₂₀N₃O: 318.1606. Found: 318.1606. Anal. (C₂₀H₂₁N₃O₂·3.5 H₂O) C, H, N.


Example YYYY:**1,5-Dihydro-3-(4-trifluoromethylphenyl-[1,2]diazepino[4,5,6-cd]-indol-6-one**

5

In a manner similar to that described for Compound 28, a solution of methyl indole-4-carboxylate (250 mg, 1.43 mmol) in dichloroethane (3 mL) was treated with *p*-trifluoromethylbenzoyl chloride (445 mg, 2.14 mmol) and aluminum chloride (572 mg). The intermediate ketone (95 mg, 0.27 mmol) in MeOH (3 mL) and conc. HCl (0.05 mL) was treated, as described, with hydrazine hydrate (0.1 mL). The reaction was quenched at 0°C with 1 M NaOAc and the aqueous layer was adjusted to pH = 8 with 1 M NaOH. The product was isolated by extraction with CH₂Cl₂, and recrystallized (CH₂Cl₂/hexanes) to give 1,5-dihydro-3-(4-trifluoromethylphenyl-[1,2]diazepino[4,5,6-cd]indol-6-one, 30 mg (34%) as a yellow solid: ¹H NMR (300 MHz, d₆-DMSO) 7.24 (app br t, 1H), 7.29 (d, *J* = 2.8 Hz, 2H), 7.60 (m, 2H), 7.82 (m, 4H), 10.57 (s, 1H), 12.01 (s, 1H). HRMS (FAB, Mna⁺) Calcd for C₁₁H₁₀N₃O: 352.0674. Found: 352.0668.

15

Example ZZZZ:20 **1,5-Dihydro-3-pentafluoroethyl-[1,2]diazepino[4,5,6-cd]-indol-6-one**



In a manner similar to that described for Compound 28, a solution of methyl indole-4-carboxylate (351 mg, 2.01 mmol) in dichloroethane (7 mL) was treated with
 5 pentafluoropropionyl chloride (2.51 mmol) and aluminum chloride (575 mg). The intermediate ketone (50 mg, 0.16 mmol) in MeOH (2 mL) and conc. HCl (0.02 mL) was treated, as described, with hydrazine hydrate (0.1 mL). The reaction was quenched at 0 °C with 1 M NaOAc and the aqueous layer was adjusted to pH = 8 with 1 M NaOH. The product was isolated by extraction with CH₂Cl₂, and recrystallized
 10 (CH₂Cl₂/MeOH/hexanes) to give 1,5-dihydro-3-pentafluoroethyl-[1,2]diazepino[4,5,6-cd]-indol-6-one, 15 mg (28%) as a yellow solid: ¹H NMR (300 MHz, d₆-DMSO) 7.16 (app br t, 1H), 7.54 (m, 2H), 7.65 (m, 1H), 10.87 (s, 1H), 12.15 (s, 1H). HRMS (FAB, Mna⁺) Calcd for C₁₁H₁₀N₃O: 352.0674. Found: 352.0668.

15

PARP Enzyme Inhibition Assay:

The PARP enzyme-inhibiting activities of the compounds of the invention were assayed as described by Simonin et al. (*J. Biol. Chem.* (1993), 268:8529-8535) and Marsischky et al. (*J. Biol. Chem.* (1995), 270:3247-3254) with minor
 20 modifications as follows. Samples (50 µL) containing 20 nM purified PARP protein, 10 µg/mL DNase I-activated calf thymus DNA (sigma), 500 µM NAD⁺, 0.5 µCi [³²P]NAD⁺, 2% DMSO, and various concentrations of test compounds were incubated in sample buffer (50 mM Tris pH 8.0, 10 mM MgCl₂, 1 mM tris(carboxyethyl)phosphine·HCl) at 25°C for 5 minutes. Under these conditions, the
 25 reaction rate was linear for times up to 10 minutes. The reaction was stopped by the addition of an equal volume of ice-cold 40% trichloroacetic acid to the samples,

which were then incubated on ice for 15 minutes. The samples were then transferred to a Bio-Dot microfiltration apparatus (BioRad), filtered through Whatman GF/C glass-fiber filter paper, washed 3 times with 150 μ L of wash buffer (5% trichloroacetic acid, 1% inorganic pyrophosphate), and dried. [32 P]ADP-Ribose incorporation into the acid-insoluble material was quantitated using a PhosphorImager (Molecular Dynamics) and ImageQuant software. Inhibition constants (K_i) were calculated by non-linear regression analyses using the velocity equation for competitive inhibition (Segel, *Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems*, John Wiley & Sons, Inc., New York (1975), 100-125). In the case of tight-binding inhibitors, 5 nM enzyme was used and the reaction was incubated at 25°C for 25 minutes. K_i values for tight-binding inhibitors were calculated using the equation described by Sculley et al. (*Biochim. Biophys. Acta* (1986), 874:44-53).

15 Cytotoxicity Potentiation Assay:

A549 cells (ATCC, Rockville, MD) were seeded into 96-well cell culture plates (Falcon brand, Fisher Scientific, Pittsburgh, PA) 16 to 24 hours before experimental manipulation. Cells were then treated with a test compound (or a combination of test compounds where indicated) for either 3 days or 5 days, at a concentration of 0.4 μ M. At the end of treatments, relative cell number was determined either by MTT assay or SRB assay. For the MTT assay, 0.2 μ g/ μ l of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Sigma Chemical Co., St. Louis, MO) was added to each well of a plate, and the plate was incubated in a cell-culture incubator for 4 hours. Metabolized MTT in each well was solubilized in 150 μ l of DMSO (Sigma Chemical Co.) with shaking and quantified with a Wallac 1420 Victor plate reader (EG&G Wallac, Gaithersburg, MD) at 540 nm. For the SRB assay, cells were fixed with 10% trichloroacetic acid (Sigma Chemical Co) for an hour at 4°C. After extensively washing, fixed cells were stained for 30 minutes with 0.4% sulforhodamine B (SRB, Sigma Chemical Co.) in 1% acetic acid (Sigma Chemical Co). Unbound SRB was washed away with 1% acetic acid.

Then the cultures were air-dried, and bound dye was solubilized with 10 mM unbuffered Tris base (Sigma Chemical Co) with shaking. The bound dye was measured photometrically with the Wallac Victor plate reader at 515 nm. The ratio of the OD (optical density) value of a compound-treated culture to the OD value of a mock-treated culture, expressed in percentage, was used to quantify the cytotoxicity of a compound. The concentration at which a compound causes 50% cytotoxicity is referred to as IC_{50} . To quantify the potentiation of the cytotoxicity of topotecan or temozolomide by test compounds, a dimensionless parameter PF_{50} is used and is defined as the ratio of the IC_{50} of topotecan or temozolomide alone to the IC_{50} of topotecan or temozolomide in combination with a test compound. For the compounds of the invention, PF_{50} values were determined by testing with topotecan.

Inhibition constants (K_i values) and cytotoxicity potentiation parameters (PF_{50} values) as determined for exemplary compounds of the invention are presented in Table 1 below. If there are two K_i values for a single compound, it means that the compound K_i was tested twice.

TABLE 1.		
PARP Enzyme Inhibition and Cytotoxicity Potentiation		
Compound No.	Inhibition Constant K_i (nM)	Cytotoxicity Potentiation PF_{50}
	69	1.1
3	2.8	N.D.
6	0.7, 1	2.2
10	38	N.D.
12	4.2	1.8
13	6.2, 4.5	N.D.
14	1.4	N.D.
16	5.0	1.9
17	6.5	N.D.
18	>>1,000	N.D.
19	62	N.D.

TABLE 1. PARP Enzyme Inhibition and Cytotoxicity Potentiation		
Compound No.	Inhibition Constant K_i (nM)	Cytotoxicity Potentiation PF_{50}
20	45	N.D.
21	5.0	2.4
22	7.2	2.3
23	4.8, 3.1	2.3
24	57	N.D.
25	4.0	N.D.
26	22, 18	N.D.
27	3.4	1.3
28	4, 3.8	1
29	8	1
30	6.3	2.4
31	5	N.D.
32	11.3	N.D.
33	230	N.D.
34	3.9	N.D.
35	3.8, 5.8	N.D.
36	29	N.D.
37	24	N.D.
38	8.4	N.D.
39	4.8	N.D.
40	5.2	N.D.
41	5.1	N.D.
42	5.1	N.D.
11	7.3	N.D.
43	2.6	N.D.
OO	4.1	2.4
PP	5.3	2.3

TABLE 1: PARP Enzyme Inhibition and Cytotoxicity Potentiation		
Compound No.	Inhibition Constant K_i (nM)	Cytotoxicity Potentiation PF_{50}
QQ	5.5, 4.5	N.D.
RR	6.9	N.D.
SS	14	N.D.
TT	12.2, 4.2	N.D.
UU	10	1.8
VV	10	2.0
WW	4.4	N.D.
XX	4.6	N.D.
YY	15.1	N.D.
ZZ	9.7	N.D.
AAA	11.4	N.D.
BBB	20	N.D.
CCC	7.3	N.D.
DDD	23	N.D.
EEE	10.6	N.D.
FFF	125	N.D.
GGG	4.1	1.9
HHH	6.6	N.D.
III	40	N.D.
JJJ	5.3	N.D.
KKK	222	N.D.
LLL	32	N.D.
MMM	9.4	2.3
NNN	172	N.D.
OOO	14	N.D.
PPP	9.4	2.1
QQQ	10.2	2.3

TABLE 1.		
PARP Enzyme Inhibition and Cytotoxicity Potentiation		
Compound No.	Inhibition Constant K_i (nM)	Cytotoxicity Potentiation PF_{50}
RRR	23	N.D.
SSS	66	N.D.
TTT	26	N.D.
UUU	11.4	N.D.
VVV	9.1	N.D.
WWW	263	N.D.
XXX	370	N.D.
YYY	6.3	1.5
ZZZ	0.7	N.D.
AAAA	1.1	N.D.
BBBB	4.8	N.D.
CCCC	4.8	N.D.
DDDD	7.7	N.D.
EEEE	2.9	N.D.
FFFF	4.7	N.D.
GGGG	6.2	N.D.
HHHH	2.2	1.9
IIII	1.4	2.6
JJJJ	4.4	2.4
KKKK	9.6	N.D.
LLLL	8.6	N.D.
MMMM	16	N.D.
NNNN	10	N.D.
OOOO	13	N.D.
PPPP	32	N.D.
QQQQ	21	N.D.
RRRR	61	N.D.


TABLE 1. PARP Enzyme Inhibition and Cytotoxicity Potentiation		
Compound No.	Inhibition Constant K_i (nM)	Cytotoxicity Potentiation PF_{50}
SSSS	19	N.D.
TTTT	7.4	1.6
UUUU	5.6	2.0
VVVV	13.2	2.1
WWWW	5.7	N.D.
XXXX	18	1.7
YYYY	9	N.D.
ZZZZ	40	N.D.


Note: N.D. = not determined.


While the invention has been described by reference to preferred embodiments and specific examples, those skilled in the art will recognize that various changes and modifications can be made without departing from the spirit and scope of the invention. Thus, the invention should be understood as not being limited by the foregoing detailed description, but as being defined by the appended claims and their equivalents.

WHAT IS CLAIMED IS:

1. A compound selected from the group consisting of:







or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof.


2. A compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 1, having a PARP-inhibiting activity corresponding to a K_i of 100 μM or less in a PARP enzyme inhibition assay.


3. A compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 1, having a cytotoxicity potentiation activity corresponding to a PF_{50} of at least 1 in a cytotoxicity potentiation assay.


4. A pharmaceutical composition comprising:


(a) an effective amount of a PARP-inhibiting agent that is:

(i) a compound selected from the group consisting of:









(ii) a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof; and


(b) a pharmaceutically acceptable carrier for said PARP-inhibiting agent.

5. A method of inhibiting PARP activity of an enzyme, comprising contacting the enzyme with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 1.

6. A method according to claim 5, wherein the enzyme is poly(ADP-ribose) polymerase or tankyrase.

7. A method of inhibiting PARP enzyme activity in mammalian tissue by administering to a mammal a therapeutically effective amount of a compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 1.

8. A method of inhibiting PARP activity of an enzyme, comprising contacting the enzyme with an effective amount of a compound of the formula:



wherein:

R^1 is: H;

halogen;

cyano;

an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; or

$-C(O)-R^{10}$, where R^{10} is: H; an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; or OR^{100} or $NR^{100}R^{110}$, where R^{100} and R^{110} are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group

R^2 is H or alkyl;

R^3 is H or alkyl;

R^4 is H, halogen or alkyl;

X is O or S;

~~X~~ Y is $(CR^5R^6)(CR^7R^8)_n$ or $N=C(R^5)$, where:


n is 0 or 1;

R^5 and R^6 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; and


R^7 and R^8 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group;

or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate of the compound.

why
NO 4 = C-R
look at #12 et.



9. A compound of the formula:



wherein:

R^1 is: H;
 halogen;
 cyano;
 an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; or
 $-C(O)-R^{10}$, where R^{10} is: H; an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; or OR^{100} or $NR^{100}R^{110}$, where R^{100} and R^{110} are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group

R^2 is H or alkyl;

R^3 is H or alkyl;

R^4 is H, halogen or alkyl;

X is O or S;

Y is $(CR^5R^6)(CR^7R^8)_n$ or $N=C(R^5)$, where:
 n is 0 or 1;


R^5 and R^6 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; and

R^7 and R^8 are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group;

where when R^1 , R^4 , R^5 , R^6 , and R^7 are each H, R^8 is not unsubstituted phenyl;

or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate of the compound.

10. A compound of the formula:



wherein:

p is 1 or 2;

R^{11} is H or alkyl;

R^{12} is halogen or an optionally substituted aryl, alkyl, alkenyl, alkynyl or acyl group $-C(O)-R^{10}$ where R^{10} is: H; an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group; or OR^{100} or $NR^{100}R^{110}$, where R^{100} and R^{110} are each independently H or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group

R^{13} is H or alkyl; and


R^{14} is H or halogen;

or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate of the compound.

11. A compound of the formula




Detailed Desc. of the Invention
- Preferred Embodiments

R¹⁵ = Me, Ph




claim

R¹⁵ = Me



(III)



wherein:

R¹⁵ is H, halogen, or an alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and alkyl and aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino;




R¹⁶ is H; halogen; cyano; or an alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, amino, and alkyl and aryl groups unsubstituted or substituted with one or more substituents selected from halogen, hydroxy, nitro, and amino;











R¹⁷ is H or alkyl; and










R¹⁸ is H, halogen, or alkyl;











where R¹⁵, R¹⁶, R¹⁷ and R¹⁸ are not all H.











12. A compound selected from the group consisting of






































wrong place?
look at p.121
claim #4



or a

pharmaceutically acceptable salt, prodrug, active metabolite or solvate thereof.

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 00/00411

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 (C07D519/00,487:00,487:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 88, no. 17, 1978 Columbus, Ohio, US; abstract no. 121125v, C.V. ANANTHANARAYANAN ET AL.: "3,4-Bridged indoles: Part II. Synthesis of 6-keto-1,5-dihydro-4,5-diazepino[6,5,4-cd]i ndoles and 3,4-disubstituted indoles as 5-HT antagonists" page 543; XP002138623 compounds of formula I & INDIAN J. CHEM., SECT B, 1997, 15(8), 710-14, -/-	9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 May 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 525 584 A (KALI CHEMIE PHARMA GMBH) 3 February 1993 (1993-02-03) examples 1-9, 11-17, 19-35, 37-50	9
X	CHEMICAL ABSTRACTS, vol. 111, no. 11, 1989 Columbus, Ohio, US; abstract no. 97216w, M. SOMEI ET AL.: "Azepinoindole derivatives as ergoline alkaloid-type pharmaceuticals" page 743; XP002138624 abstract -& DATABASE CHEMICAL ABSTRACTS 'Online! Chemical Abstracts Service, Columbus, Ohio, US 111:97216w, XP002138625 compound with RN '122328-80-7! abstract & JP 00 134988 A (KISSEI) 6 February 1989 (1989-02-06)	9
X	R.E. BOWMAN ET AL.: "1,3,4,5-tetrahydrobenz[cd]indoles and related compounds. Part II" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1., 1972, pages 1926-32, XP002138621 CHEMICAL SOCIETY, LETCHWORTH., GB ISSN: 0300-922X compounds 37, 38 and 39	9
A	WO 97 04771 A (UNIVERSITY NEWCASTLE VENTURES LTD) 13 February 1997 (1997-02-13) claim 1	4, 15
P, X	WO 99 59975 A (GUILFORD PHARM INC) 25 November 1999 (1999-11-25) claims 1, 10, 14, 23, 79	1, 4, 5, 9

13. A compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 9, having a PARP-inhibiting activity corresponding to a K_i of 100 μ M or less in a PARP enzyme inhibition assay.

14. A compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 9, having a cytotoxicity potentiation activity corresponding to a PF_{50} of at least 1 in a cytotoxicity potentiation assay.

15. A pharmaceutical composition comprising:

- (a) an effective amount of a PARP-inhibiting agent that is a compound according to claim 9; or a pharmaceutically acceptable salt, prodrug, active metabolite, or solvate thereof; and
- (b) a pharmaceutically acceptable carrier for said PARP-inhibiting agent.

16. A method of inhibiting PARP activity of an enzyme, comprising contacting the enzyme with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 9.

17. A method according to claim 16, wherein the enzyme is poly(ADP-ribose) polymerase or tankyrase.

18. A method of inhibiting PARP enzyme activity in mammalian tissue by administering to a mammal a therapeutically effective amount of a compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 9.

19. A method of inhibiting PARP activity of an enzyme, comprising contacting the enzyme with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, active metabolite, or solvate according to claim 12.

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/US 00/00411

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/06 C07D487/06 A61K31/4375 A61K31/55 A61K31/551
 A61P35/00 C07D519/00 //(C07D471/06,221:00,209:00),
 (C07D487/06,223:00,209:00),(C07D487/06,243:00,209:00),

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 75, no. 23, 1971 Columbus, Ohio, US; abstract no. 140614, MUCHOWSKI ET AL.: "Isocarbostyrils. II. Conversion of 2-methyl-4-acyl-5-nitroisocarbostyrils to 2-substituted indole-4-carboxylic acids" page 304; XP002138622 compounds of formula II & CAN. J. CHEM., vol. 49, no. 17, 1971, pages 2797-802,	9,10
X	US 3 900 477 A (PHILIPP ET AL) 19 August 1975 (1975-08-19) example 4	9
	--- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 May 2000

Date of mailing of the international search report

28/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I